

Table 3A
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic

Page: 1
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Racial and Ethnic Distribution ⁽²⁾							
		White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %	
0101	60	*	*	*	*	*	*	*	
0102	180	95.6	1.1	1.7	0.6	0.0	1.1	0.0	
0103	183	97.3	0.0	1.1	0.0	0.5	1.1	0.0	
0104	54	*	*	*	*	*	*	*	
0250	60	*	*	*	*	*	*	*	
0251	63	*	*	*	*	*	*	*	
0252	51	*	*	*	*	*	*	*	
0253	72	*	*	*	*	*	*	*	
0401	235	94.5	0.9	1.3	0.0	0.4	2.1	0.9	
0402	169	84.0	0.0	0.6	0.0	1.8	13.6	0.0	
0403	57	*	*	*	*	*	*	*	
0404	56	*	*	*	*	*	*	*	
0501	225	94.2	0.0	2.2	0.0	0.4	3.1	0.0	
0502	1	*	*	*	*	*	*	*	
0503	79	*	*	*	*	*	*	*	
0504	232	91.4	0.0	0.9	1.7	0.9	5.2	0.0	
0505	145	86.2	2.1	3.4	0.7	0.0	7.6	0.0	
0506	75	*	*	*	*	*	*	*	
0650	479	90.8	3.5	1.7	1.0	1.0	1.5	0.4	
0651	268	94.0	0.4	1.9	0.4	0.0	2.2	1.1	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Racial and Ethnic Distribution by Clinic

Page: 2
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Racial and Ethnic Distribution (2)							
		White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %	
0652	124	96.8	0.0	0.8	0.0	0.0	0.8	1.6	
0653	173	98.8	0.0	0.6	0.0	0.0	0.6	0.0	
0654	168	92.3	1.2	1.8	1.2	0.0	3.0	0.6	
0655	153	96.7	0.0	0.0	0.0	0.0	3.3	0.0	
0657	29	*	*	*	*	*	*	*	
0701	173	91.9	0.6	1.2	0.0	5.2	0.6	0.6	
0702	30	*	*	*	*	*	*	*	
0703	100	98.0	0.0	1.0	1.0	0.0	0.0	0.0	
0704	26	*	*	*	*	*	*	*	
0705	64	*	*	*	*	*	*	*	
0850	486	89.7	3.5	4.1	0.0	1.0	1.2	0.4	
0851	359	94.2	0.8	3.6	0.0	0.3	0.6	0.6	
0901	722	84.2	2.1	8.9	0.1	0.1	4.3	0.3	
0902	28	*	*	*	*	*	*	*	
1051	595	90.6	1.0	4.2	0.7	0.2	2.4	1.0	
1101	835	37.7	53.2	4.6	0.2	0.8	1.6	1.9	
1102	371	81.1	10.8	5.1	0.0	0.8	1.6	0.5	
1106	46	*	*	*	*	*	*	*	
1250	446	86.5	0.2	9.6	0.0	1.6	1.8	0.2	
1251	353	92.9	0.8	2.5	0.0	0.0	1.1	2.5	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic

Page: 3
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Racial and Ethnic Distribution ⁽²⁾							
		White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %	
1252	253	65.2	0.8	32.4	0.0	0.4	0.8	0.4	
1253	381	82.7	4.7	8.1	0.0	0.0	3.4	1.0	
1350	1,025	63.8	20.3	8.4	0.2	2.5	3.5	1.3	
1351	71	*	*	*	*	*	*	*	
1352	167	64.1	25.7	5.4	0.0	0.6	3.6	0.6	
1401	95	*	*	*	*	*	*	*	
1402	73	*	*	*	*	*	*	*	
1403	9	*	*	*	*	*	*	*	
1404	12	*	*	*	*	*	*	*	
1501	480	45.4	6.5	39.0	0.6	0.0	1.3	7.3	
1503	580	60.7	5.9	25.9	0.7	1.0	0.9	5.0	
1504	1,052	49.3	4.5	37.5	0.1	5.1	1.9	1.6	
1505	369	88.1	1.6	5.4	0.0	1.1	2.4	1.4	
1506	266	80.8	1.5	13.5	0.8	0.4	1.9	1.1	
1507	266	93.6	0.4	2.6	0.0	0.8	1.5	1.1	
1508	72	*	*	*	*	*	*	*	
1509	309	52.1	1.3	41.1	0.0	0.3	1.0	4.2	
1510	46	*	*	*	*	*	*	*	
1511	281	94.3	1.8	2.1	0.0	0.0	0.7	1.1	
1512	109	95.4	0.0	1.8	0.0	0.0	2.8	0.0	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Racial and Ethnic Distribution by Clinic

Page: 4
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Racial and Ethnic Distribution (2)							
		White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %	
1701	311	62.1	0.0	1.6	23.8	1.6	10.3	0.6	
1712	21	*	*	*	*	*	*	*	
1950	320	91.9	0.3	6.3	0.0	0.0	1.6	0.0	
1952	53	*	*	*	*	*	*	*	
1953	125	92.0	0.0	4.0	0.0	1.6	2.4	0.0	
1954	300	90.0	0.0	8.7	0.0	0.3	0.3	0.7	
1955	59	*	*	*	*	*	*	*	
1957	218	90.4	0.9	4.6	0.9	0.0	1.8	1.4	
1959	67	*	*	*	*	*	*	*	
2150	296	95.6	0.0	1.0	0.3	0.0	2.0	1.0	
2151	128	92.2	0.0	1.6	0.8	0.8	2.3	2.3	
2251	193	89.6	0.0	1.6	0.0	0.5	7.3	1.0	
2252	111	95.5	0.0	2.7	0.0	0.0	1.8	0.0	
2551	1,359	32.5	64.5	1.6	0.0	0.2	1.1	0.1	
2554	38	*	*	*	*	*	*	*	
2555	1,673	74.6	20.3	3.3	0.1	0.4	1.0	0.3	
3001	408	10.3	68.6	18.6	0.0	1.0	1.5	0.0	
3002	100	22.0	55.0	21.0	0.0	0.0	2.0	0.0	
3003	4	*	*	*	*	*	*	*	
3004	72	*	*	*	*	*	*	*	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic

Page: 5
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Racial and Ethnic Distribution ⁽²⁾							
		White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %	
3007	12	*	*	*	*	*	*	*	
3009	9	*	*	*	*	*	*	*	
3010	3	*	*	*	*	*	*	*	
3201	236	94.9	0.0	2.5	0.0	1.3	0.4	0.8	
3301	1,808	47.4	26.5	16.2	0.1	4.2	4.6	1.0	
3302	110	26.4	47.3	15.5	0.0	5.5	4.5	0.9	
3304	40	*	*	*	*	*	*	*	
3305	40	*	*	*	*	*	*	*	
3306	88	*	*	*	*	*	*	*	
3308	19	*	*	*	*	*	*	*	
3309	12	*	*	*	*	*	*	*	
3550	34	*	*	*	*	*	*	*	
3551	9	*	*	*	*	*	*	*	
3552	14	*	*	*	*	*	*	*	
3701	443	93.0	0.2	5.6	0.0	0.2	0.9	0.0	
3805	1,387	76.6	13.1	6.3	0.0	0.9	2.8	0.2	
3950	864	68.1	17.1	6.5	0.2	2.0	5.7	0.5	
3952	5	*	*	*	*	*	*	*	
3953	3	*	*	*	*	*	*	*	
3954	25	*	*	*	*	*	*	*	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic

Page: 6
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Racial and Ethnic Distribution ⁽²⁾							
		White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %	
4001	480	30.6	58.1	7.7	0.6	0.4	2.1	0.4	
4002	391	51.2	25.3	15.3	0.3	1.5	5.9	0.5	
4103	33	*	*	*	*	*	*	*	
4106	338	14.8	47.3	32.2	0.0	0.6	3.3	1.8	
4107	1,553	51.3	27.2	14.7	0.1	2.5	3.9	0.3	
4109	512	91.4	0.4	5.5	0.0	0.6	1.6	0.6	
4110	722	40.2	9.1	45.7	0.3	0.7	3.6	0.4	
4112	254	0.8	0.0	98.0	0.0	0.0	0.4	0.8	
4113	271	5.5	19.9	70.8	0.0	0.7	3.0	0.0	
4120	861	61.1	8.0	25.1	0.1	2.4	3.0	0.2	
4160	752	43.6	32.7	11.4	0.0	7.7	4.4	0.1	
4351	253	93.3	0.8	3.2	0.0	0.0	1.2	1.6	
4352	73	*	*	*	*	*	*	*	
4353	207	68.1	0.5	26.6	0.0	0.0	2.9	1.9	
4354	342	92.4	2.0	1.5	0.0	0.6	3.5	0.0	
4355	159	78.6	0.0	6.9	0.0	0.0	11.9	2.5	
4356	239	82.4	0.0	11.3	1.3	0.8	3.3	0.8	
4357	4	*	*	*	*	*	*	*	
4358	96	*	*	*	*	*	*	*	
4359	239	72.4	0.4	23.4	0.4	0.4	1.3	1.7	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic

Page: 7
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Racial and Ethnic Distribution ⁽²⁾							
		White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %	
4360	63	*	*	*	*	*	*	*	
4361	110	89.1	1.8	0.9	0.0	0.9	7.3	0.0	
4362	150	94.7	0.0	1.3	0.7	0.0	2.7	0.7	
4363	369	91.6	0.3	3.5	0.0	0.8	3.0	0.8	
4401	572	90.2	0.7	3.8	0.0	0.3	4.5	0.3	
4501	394	94.4	0.5	4.1	0.0	0.0	0.8	0.3	
4601	676	76.3	1.3	17.6	0.0	0.4	1.5	2.8	
4701	536	92.4	0.6	3.2	0.2	1.7	0.9	1.1	
5050	579	68.6	24.2	2.2	0.3	0.9	3.6	0.2	
5051	1,140	65.4	25.4	3.5	0.1	1.5	3.7	0.5	
5052	905	66.1	21.5	1.9	0.2	7.8	2.0	0.4	
5101	970	72.7	8.2	12.2	0.1	1.5	4.5	0.7	
5202	268	88.8	0.0	1.5	1.9	0.4	7.5	0.0	
5203	71	*	*	*	*	*	*	*	
5204	79	*	*	*	*	*	*	*	
5601	494	94.5	1.2	1.6	0.0	0.4	1.6	0.6	
5801	965	88.0	4.6	4.5	0.3	0.1	1.1	1.5	
5803	20	*	*	*	*	*	*	*	
5804	7	*	*	*	*	*	*	*	
6101	17	*	*	*	*	*	*	*	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic

Page: 8
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Racial and Ethnic Distribution ⁽²⁾							
		White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %	
6102	66	*	*	*	*	*	*	*	
6103	1,727	61.8	25.9	7.4	0.1	0.4	3.9	0.5	
6104	41	*	*	*	*	*	*	*	
6105	18	*	*	*	*	*	*	*	
6106	3	*	*	*	*	*	*	*	
6301	2,000	49.3	26.6	18.8	0.1	2.4	2.3	0.4	
6312	1,072	41.8	51.2	1.5	0.1	2.5	2.2	0.7	
6316	599	72.8	14.0	7.8	0.3	2.5	1.7	0.8	
6320	101	71.3	21.8	2.0	0.0	4.0	1.0	0.0	
6325	61	*	*	*	*	*	*	*	
6330	217	29.5	51.2	17.1	0.0	0.0	2.3	0.0	
6335	258	77.1	17.4	1.9	0.0	2.7	0.8	0.0	
6340	60	*	*	*	*	*	*	*	
6345	22	*	*	*	*	*	*	*	
6350	115	57.4	37.4	3.5	0.0	0.9	0.9	0.0	
7304	95	*	*	*	*	*	*	*	
7306	1,102	51.0	33.4	12.3	0.1	0.4	2.2	0.6	
7308	80	*	*	*	*	*	*	*	
7310	26	*	*	*	*	*	*	*	
7401	918	85.7	4.6	5.2	0.0	0.7	3.6	0.2	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic

Page: 9
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Racial and Ethnic Distribution ⁽²⁾							
		White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %	
7402	56	*	*	*	*	*	*	*	
7403	28	*	*	*	*	*	*	*	
7404	26	*	*	*	*	*	*	*	
7601	301	94.0	0.0	4.0	0.0	0.0	0.7	1.3	
7850	391	91.0	0.0	5.4	0.0	0.0	2.3	1.3	
7851	54	*	*	*	*	*	*	*	
7852	39	*	*	*	*	*	*	*	
8150	20	*	*	*	*	*	*	*	
8151	1,230	43.7	41.1	8.7	0.6	2.4	3.4	0.1	
8152	200	37.5	50.0	1.5	0.5	1.5	9.0	0.0	
8153	10	*	*	*	*	*	*	*	
8154	106	34.9	17.9	12.3	0.0	34.0	0.9	0.0	
8155	22	*	*	*	*	*	*	*	
8156	10	*	*	*	*	*	*	*	
8157	14	*	*	*	*	*	*	*	
8201	1,025	60.9	26.7	5.1	0.3	2.6	4.4	0.0	
8202	1,233	89.2	5.7	2.5	0.2	0.9	1.3	0.2	
8203	137	73.7	19.7	5.1	0.0	0.0	0.7	0.7	
8250	345	44.9	46.7	2.9	0.0	2.9	2.6	0.0	
8255	996	70.7	21.0	4.8	0.0	1.9	1.4	0.2	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic

Page: 10
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Racial and Ethnic Distribution ⁽²⁾							
		White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %	
8257	221	46.2	28.1	0.0	0.0	24.4	1.4	0.0	
8258	608	63.8	27.8	5.1	0.0	0.7	2.1	0.5	
8260	184	46.7	34.8	1.6	0.0	14.7	2.2	0.0	
8901	375	0.5	94.9	1.1	0.0	0.8	2.7	0.0	
8902	259	2.7	95.0	1.2	0.0	0.0	1.2	0.0	
8903	478	2.5	95.0	1.3	0.0	0.2	1.0	0.0	
8904	352	4.8	81.3	0.9	0.0	9.4	3.7	0.0	
8905	1,124	36.7	30.1	24.1	0.1	5.3	3.4	0.4	
9004	4	*	*	*	*	*	*	*	
9012	15	*	*	*	*	*	*	*	
9013	30	*	*	*	*	*	*	*	
9015	10	*	*	*	*	*	*	*	
9016	34	*	*	*	*	*	*	*	
9017	10	*	*	*	*	*	*	*	
9018	33	*	*	*	*	*	*	*	
9019	14	*	*	*	*	*	*	*	
9021	20	*	*	*	*	*	*	*	
9040	7	*	*	*	*	*	*	*	
9041	180	2.2	95.6	1.1	0.0	0.0	1.1	0.0	
9045	502	9.8	45.4	42.6	0.0	0.2	1.6	0.4	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2009 Pregnancy Nutrition Surveillance ⁽¹⁾
Michigan
Comparison of Racial and Ethnic Distribution by Clinic

Page: 11
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Racial and Ethnic Distribution ⁽²⁾							
		White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %	
9050	811	13.6	58.7	26.0	0.0	0.9	0.6	0.2	
9051	962	8.8	12.6	77.8	0.0	0.2	0.6	0.0	
9053	1,113	2.2	96.0	0.6	0.0	0.5	0.5	0.2	
9055	52	*	*	*	*	*	*	*	
9058	260	73.5	19.6	0.4	0.0	5.8	0.8	0.0	
9066	945	1.1	98.2	0.1	0.2	0.2	0.2	0.0	
9068	572	2.3	95.5	0.9	0.0	0.3	0.5	0.5	
9069	1,104	2.1	96.2	0.5	0.0	0.2	1.0	0.1	
9070	831	1.6	89.3	0.1	0.2	7.2	1.6	0.0	
9072	731	1.5	94.7	0.3	0.3	2.6	0.4	0.3	
9074	705	3.1	94.6	0.6	0.0	0.1	1.3	0.3	
9080	921	23.0	72.1	1.8	0.0	0.2	2.3	0.5	
Unknown Clinics	52	*	*	*	*	*	*	*	
Michigan	69,465	57.1	27.2	10.5	0.3	1.7	2.5	0.7	
Nation (Prior Year)	1,307,887	42.0	24.2	28.1	1.0	3.1	1.1	0.5	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Age Distribution by Clinic

Page: 12
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Age Distribution ⁽²⁾							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
0101	60	*	*	*	*	*	*	*	*
0102	180	0.6	5.0	13.3	65.6	14.4	1.1	0.0	
0103	183	0.0	3.8	12.0	66.7	14.8	2.7	0.0	
0104	54	*	*	*	*	*	*	*	*
0250	60	*	*	*	*	*	*	*	*
0251	63	*	*	*	*	*	*	*	*
0252	51	*	*	*	*	*	*	*	*
0253	72	*	*	*	*	*	*	*	*
0401	235	0.4	6.8	17.9	64.3	10.6	0.0	0.0	
0402	169	0.0	4.1	13.0	59.2	22.5	1.2	0.0	
0403	57	*	*	*	*	*	*	*	*
0404	56	*	*	*	*	*	*	*	*
0501	225	0.0	4.0	16.0	63.6	15.6	0.9	0.0	
0502	1	*	*	*	*	*	*	*	*
0503	79	*	*	*	*	*	*	*	*
0504	232	0.0	3.9	15.5	61.2	18.5	0.9	0.0	
0505	145	0.0	2.1	12.4	68.3	16.6	0.7	0.0	
0506	75	*	*	*	*	*	*	*	*
0650	479	0.0	5.2	12.9	65.3	15.4	0.8	0.2	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Age Distribution by Clinic

Page: 13
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Age Distribution ⁽²⁾							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
0651	268	0.4	6.0	13.1	67.9	12.3	0.4	0.0	
0652	124	0.0	7.3	11.3	62.1	14.5	4.0	0.8	
0653	173	1.2	5.2	13.9	66.5	10.4	2.3	0.6	
0654	168	0.0	7.7	13.1	61.3	16.7	1.2	0.0	
0655	153	0.0	5.9	15.7	59.5	19.0	0.0	0.0	
0657	29	*	*	*	*	*	*	*	*
0701	173	0.0	2.9	8.1	67.6	19.1	2.3	0.0	
0702	30	*	*	*	*	*	*	*	*
0703	100	0.0	1.0	12.0	66.0	20.0	1.0	0.0	
0704	26	*	*	*	*	*	*	*	*
0705	64	*	*	*	*	*	*	*	*
0850	486	0.2	5.1	10.5	67.3	16.5	0.4	0.0	
0851	359	0.0	6.7	14.5	66.6	11.1	1.1	0.0	
0901	722	0.3	6.9	13.7	60.7	17.0	1.2	0.1	
0902	28	*	*	*	*	*	*	*	*
1051	595	0.5	4.9	11.4	62.4	19.0	1.8	0.0	
1101	835	0.1	10.3	14.5	58.8	14.5	1.7	0.1	
1102	371	0.5	5.7	14.0	63.3	15.1	1.1	0.3	
1106	46	*	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Age Distribution by Clinic

Page: 14
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Age Distribution ⁽²⁾							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
1250	446	0.4	5.6	11.2	68.6	13.7	0.4	0.0	
1251	353	0.3	5.4	12.2	66.6	15.0	0.6	0.0	
1252	253	0.4	4.3	12.3	60.9	20.2	2.0	0.0	
1253	381	0.5	4.5	15.2	64.0	15.0	0.8	0.0	
1350	1,025	0.3	7.3	14.3	62.0	15.2	0.7	0.1	
1351	71	*	*	*	*	*	*	*	*
1352	167	0.0	6.6	13.2	64.7	13.2	2.4	0.0	
1401	95	*	*	*	*	*	*	*	*
1402	73	*	*	*	*	*	*	*	*
1403	9	*	*	*	*	*	*	*	*
1404	12	*	*	*	*	*	*	*	*
1501	480	0.4	8.1	13.8	59.0	17.7	1.0	0.0	
1503	580	0.0	7.6	12.9	59.3	18.4	1.4	0.3	
1504	1,052	0.5	6.9	10.5	61.0	20.2	0.9	0.0	
1505	369	0.0	3.5	10.0	60.4	24.9	1.1	0.0	
1506	266	0.0	7.9	13.2	60.2	17.7	1.1	0.0	
1507	266	0.0	3.4	10.2	66.5	19.2	0.8	0.0	
1508	72	*	*	*	*	*	*	*	*
1509	309	0.3	7.8	13.3	58.6	17.8	2.3	0.0	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Age Distribution by Clinic

Page: 15
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Age Distribution (2)							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
1510	46	*	*	*	*	*	*	*	*
1511	281	0.0	7.1	12.1	66.9	12.1	1.8	0.0	
1512	109	0.0	2.8	10.1	59.6	24.8	2.8	0.0	
1701	311	0.0	4.8	13.2	65.0	17.0	0.0	0.0	
1712	21	*	*	*	*	*	*	*	*
1950	320	0.0	7.2	20.6	60.0	10.0	0.9	1.3	
1952	53	*	*	*	*	*	*	*	*
1953	125	0.0	2.4	17.6	66.4	12.8	0.8	0.0	
1954	300	0.3	3.7	15.7	67.3	12.3	0.7	0.0	
1955	59	*	*	*	*	*	*	*	*
1957	218	0.0	5.5	11.0	61.0	21.1	1.4	0.0	
1959	67	*	*	*	*	*	*	*	*
2150	296	0.0	3.7	8.1	67.9	19.6	0.7	0.0	
2151	128	0.8	6.3	7.0	64.8	18.8	1.6	0.8	
2251	193	0.5	5.2	11.4	65.3	16.6	1.0	0.0	
2252	111	2.7	7.2	11.7	64.9	13.5	0.0	0.0	
2551	1,359	0.5	9.9	19.8	56.4	12.0	1.2	0.1	
2554	38	*	*	*	*	*	*	*	*
2555	1,673	0.1	5.9	12.7	61.6	18.5	1.1	0.1	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Age Distribution by Clinic

Page: 16
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Age Distribution ⁽²⁾							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
3001	408	0.7	12.3	18.6	53.2	13.2	1.7	0.2	
3002	100	0.0	9.0	9.0	63.0	19.0	0.0	0.0	
3003	4	*	*	*	*	*	*	*	*
3004	72	*	*	*	*	*	*	*	*
3007	12	*	*	*	*	*	*	*	*
3009	9	*	*	*	*	*	*	*	*
3010	3	*	*	*	*	*	*	*	*
3201	236	0.0	3.0	11.9	62.3	20.3	2.5	0.0	
3301	1,808	0.3	6.4	13.6	60.5	18.0	1.3	0.0	
3302	110	0.0	4.5	14.5	60.9	19.1	0.9	0.0	
3304	40	*	*	*	*	*	*	*	*
3305	40	*	*	*	*	*	*	*	*
3306	88	*	*	*	*	*	*	*	*
3308	19	*	*	*	*	*	*	*	*
3309	12	*	*	*	*	*	*	*	*
3550	34	*	*	*	*	*	*	*	*
3551	9	*	*	*	*	*	*	*	*
3552	14	*	*	*	*	*	*	*	*
3701	443	0.0	4.7	12.2	71.1	10.8	1.1	0.0	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Age Distribution by Clinic

Page: 17
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Age Distribution (2)							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
3805	1,387	0.5	6.9	15.0	61.2	13.6	1.4	1.4	
3950	864	0.5	5.4	12.7	63.7	17.1	0.6	0.0	
3952	5	*	*	*	*	*	*	*	*
3953	3	*	*	*	*	*	*	*	*
3954	25	*	*	*	*	*	*	*	*
4001	480	0.6	10.4	16.0	57.3	13.5	2.1	0.0	
4002	391	0.3	6.1	13.3	64.2	14.6	1.5	0.0	
4103	33	*	*	*	*	*	*	*	*
4106	338	0.0	4.1	11.5	65.7	15.7	2.1	0.9	
4107	1,553	0.3	8.0	15.3	57.5	16.4	1.0	1.5	
4109	512	0.2	5.5	9.4	62.5	20.9	1.4	0.2	
4110	722	0.4	6.4	14.4	55.8	21.9	1.0	0.1	
4112	254	0.0	5.9	7.5	56.7	27.6	2.0	0.4	
4113	271	0.7	5.5	14.8	55.4	22.1	0.4	1.1	
4120	861	0.1	5.5	10.8	64.9	17.5	1.0	0.1	
4160	752	0.3	4.1	10.1	62.9	21.1	1.2	0.3	
4351	253	0.0	9.1	15.8	55.7	18.6	0.8	0.0	
4352	73	*	*	*	*	*	*	*	*
4353	207	0.0	5.8	14.5	65.2	13.0	1.4	0.0	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Age Distribution by Clinic

Page: 18
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Age Distribution ⁽²⁾							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
4354	342	0.0	6.4	17.8	63.2	11.1	1.2	0.3	
4355	159	0.0	5.0	13.8	66.0	15.1	0.0	0.0	
4356	239	0.0	7.1	17.6	62.3	12.1	0.8	0.0	
4357	4	*	*	*	*	*	*	*	*
4358	96	*	*	*	*	*	*	*	*
4359	239	0.0	4.2	14.2	60.7	20.1	0.8	0.0	
4360	63	*	*	*	*	*	*	*	*
4361	110	3.6	8.2	16.4	60.0	10.9	0.9	0.0	
4362	150	0.0	4.0	14.0	69.3	12.7	0.0	0.0	
4363	369	0.0	5.1	10.0	69.6	14.4	0.8	0.0	
4401	572	0.0	3.7	14.9	62.9	17.7	0.9	0.0	
4501	394	0.3	4.3	10.7	64.0	19.8	0.8	0.3	
4601	676	0.0	5.2	14.9	61.8	17.0	1.0	0.0	
4701	536	0.0	3.7	10.6	62.1	22.4	1.1	0.0	
5050	579	0.2	4.3	11.2	58.5	24.0	1.7	0.0	
5051	1,140	0.2	3.5	9.3	63.2	22.0	1.6	0.2	
5052	905	0.2	4.5	9.7	58.1	26.0	1.4	0.0	
5101	970	0.0	3.2	10.1	59.2	24.8	2.6	0.1	
5202	268	0.0	1.9	11.9	66.4	19.4	0.4	0.0	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Age Distribution by Clinic

Page: 19
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Age Distribution ⁽²⁾							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
5203	71	*	*	*	*	*	*	*	*
5204	79	*	*	*	*	*	*	*	*
5601	494	0.0	4.0	12.3	68.2	14.8	0.6	0.0	
5801	965	0.2	6.8	14.0	62.0	16.0	1.0	0.0	
5803	20	*	*	*	*	*	*	*	*
5804	7	*	*	*	*	*	*	*	*
6101	17	*	*	*	*	*	*	*	*
6102	66	*	*	*	*	*	*	*	*
6103	1,727	0.2	6.8	15.2	62.8	14.2	0.8	0.0	
6104	41	*	*	*	*	*	*	*	*
6105	18	*	*	*	*	*	*	*	*
6106	3	*	*	*	*	*	*	*	*
6301	2,000	0.3	5.8	12.2	60.4	20.0	1.3	0.0	
6312	1,072	0.3	2.7	8.6	57.2	28.4	2.8	0.1	
6316	599	0.2	3.3	6.8	63.8	24.0	1.8	0.0	
6320	101	0.0	4.0	8.9	58.4	27.7	1.0	0.0	
6325	61	*	*	*	*	*	*	*	*
6330	217	0.9	9.7	24.0	46.5	18.4	0.5	0.0	
6335	258	0.0	3.1	6.6	61.6	25.6	3.1	0.0	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Age Distribution by Clinic

Page: 20
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Age Distribution (2)							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
6340	60	*	*	*	*	*	*	*	*
6345	22	*	*	*	*	*	*	*	*
6350	115	0.0	7.8	11.3	57.4	20.9	1.7	0.9	
7304	95	*	*	*	*	*	*	*	*
7306	1,102	0.4	7.8	14.8	57.9	17.8	1.2	0.2	
7308	80	*	*	*	*	*	*	*	*
7310	26	*	*	*	*	*	*	*	*
7401	918	0.3	5.6	13.6	65.6	13.9	1.0	0.0	
7402	56	*	*	*	*	*	*	*	*
7403	28	*	*	*	*	*	*	*	*
7404	26	*	*	*	*	*	*	*	*
7601	301	0.3	4.3	9.0	68.4	16.3	1.3	0.3	
7850	391	0.3	6.1	14.6	60.9	16.1	2.0	0.0	
7851	54	*	*	*	*	*	*	*	*
7852	39	*	*	*	*	*	*	*	*
8150	20	*	*	*	*	*	*	*	*
8151	1,230	0.0	3.3	8.1	64.5	22.3	1.7	0.1	
8152	200	0.5	15.5	33.0	50.5	0.0	0.0	0.5	
8153	10	*	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Age Distribution by Clinic

Page: 21
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Age Distribution ⁽²⁾							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
8154	106	0.0	0.9	3.8	56.6	37.7	0.9	0.0	
8155	22	*	*	*	*	*	*	*	*
8156	10	*	*	*	*	*	*	*	*
8157	14	*	*	*	*	*	*	*	*
8201	1,025	0.1	5.1	13.2	57.3	22.8	1.5	0.1	
8202	1,233	0.1	2.8	7.7	60.7	25.1	3.3	0.3	
8203	137	0.0	4.4	13.1	62.0	16.8	3.6	0.0	
8250	345	0.0	4.1	15.4	59.1	20.0	1.4	0.0	
8255	996	0.3	4.5	11.7	60.5	16.9	1.0	5.0	
8257	221	0.0	3.6	10.0	63.3	20.4	2.7	0.0	
8258	608	0.3	5.3	11.2	59.0	22.4	1.8	0.0	
8260	184	0.0	4.3	10.3	60.9	20.7	2.7	1.1	
8901	375	1.6	10.1	12.8	58.4	16.5	0.5	0.0	
8902	259	0.8	3.5	16.2	59.5	18.1	1.9	0.0	
8903	478	1.3	10.9	15.5	53.1	17.6	1.7	0.0	
8904	352	0.6	6.3	11.9	55.7	24.4	0.9	0.3	
8905	1,124	0.4	6.0	10.7	56.6	24.2	1.9	0.3	
9004	4	*	*	*	*	*	*	*	*
9012	15	*	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Age Distribution by Clinic

Page: 22
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Age Distribution ⁽²⁾							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
9013	30	*	*	*	*	*	*	*	*
9015	10	*	*	*	*	*	*	*	*
9016	34	*	*	*	*	*	*	*	*
9017	10	*	*	*	*	*	*	*	*
9018	33	*	*	*	*	*	*	*	*
9019	14	*	*	*	*	*	*	*	*
9021	20	*	*	*	*	*	*	*	*
9040	7	*	*	*	*	*	*	*	*
9041	180	0.0	8.3	17.8	52.2	21.1	0.6	0.0	
9045	502	1.2	14.9	15.9	50.8	15.3	1.6	0.2	
9050	811	0.5	7.9	12.2	53.4	22.9	2.8	0.2	
9051	962	0.3	7.1	10.7	54.8	24.9	2.0	0.2	
9053	1,113	0.2	8.0	16.0	55.1	19.0	1.5	0.2	
9055	52	*	*	*	*	*	*	*	
9058	260	0.0	1.9	5.8	59.6	29.2	3.5	0.0	
9066	945	0.2	8.6	17.9	49.6	19.4	1.7	2.6	
9068	572	0.2	8.2	14.9	56.6	19.1	1.0	0.0	
9069	1,104	0.1	6.8	15.2	55.5	20.8	1.5	0.0	
9070	831	0.5	8.4	15.9	55.8	17.6	1.7	0.1	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Age Distribution by Clinic

Page: 23
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Age Distribution ⁽²⁾							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
9072	731	0.5	8.6	20.5	53.6	15.3	1.4	0.0	
9074	705	0.3	7.0	15.7	58.9	16.9	1.3	0.0	
9080	921	0.2	5.4	12.9	59.0	20.7	1.7	0.0	
Unknown Clinics	52	*	*	*	*	*	*	*	*
Michigan	69,465	0.3	6.1	13.0	60.4	18.6	1.4		0.3
Nation (Prior Year)	1,307,887	0.3	6.3	12.6	59.7	19.4	1.4		0.3

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Maternal Health Indicators by Clinic

Page: 24
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Health Indicators (2, 3)											
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Gestational Diabetes During Pregnancy	Hypertension During Pregnancy	
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester	Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0101	55	* * * *	*	49	* * * *	*	3	* *	11	* *	36	* * 36
0102	170	5.3 (93)	58.2 (103)	139	41.0 (108)	41.7 (18)	21	* *	28	* *	123	7.3 (79) 124
0103	178	2.8 (18)	54.5 (69)	155	42.6 (110)	39.4 (12)	13	* *	31	* *	117	8.5 (95) 117
0104	53	* * * *	*	47	* * * *	*	3	* *	5	* *	36	* * 36
0250	56	* * * *	*	48	* * * *	*	4	* *	13	* *	42	* * 42
0251	63	* * * *	*	53	* * * *	*	6	* *	5	* *	35	* * 35
0252	50	* * * *	*	43	* * * *	*	3	* *	14	* *	40	* * 40
0253	68	* * * *	*	60	* * * *	*	4	* *	10	* *	50	* * 50
0401	231	5.2 (89)	49.8 (24)	208	20.2 (53)	51.4 (70)	23	* *	42	* *	150	4.7 (29) 150
0402	168	3.0 (20)	50.6 (32)	149	20.1 (51)	47.7 (39)	19	* *	38	* *	131	4.6 (26) 131
0403	56	* * * *	*	47	* * * *	*	1	* *	8	* *	35	* * 35
0404	54	* * * *	*	50	* * * *	*	5	* *	13	* *	41	* * 41
0501	169	5.9 (112)	49.7 (22)	144	27.1 (90)	43.1 (21)	8	* *	14	* *	145	5.5 (46) 145
0502	1	* * * *	*	0	* * * *	*	0	* *	1	* *	1	* * 1
0503	66	* * * *	*	59	* * * *	*	5	* *	4	* *	52	* * 52
0504	170	4.1 (58)	48.8 (14)	135	23.0 (69)	45.2 (29)	14	* *	15	* *	154	3.2 (5) 154
0505	108	0.9 (2)	58.3 (104)	95	* * * *	*	14	* *	11	* *	105	8.6 (96) 106
												17.0(113)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Maternal Health Indicators by Clinic

Page: 25
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Health Indicators (2, 3)											
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Gestational Diabetes During Pregnancy	Hypertension During Pregnancy	
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester	Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0506	69	* *	* *	56	* *	* *	6	* *	4	* *	51	* *
0650	470	4.9 (82)	54.5 (69)	398	18.6 (39)	52.0 (78)	62	* *	84	* *	425	3.1 (3)
0651	257	5.4 (97)	56.4 (89)	222	18.0 (30)	51.8 (76)	27	* *	37	* *	231	3.5 (8)
0652	120	3.3 (28)	58.3 (104)	105	29.5 (100)	45.7 (30)	11	* *	30	* *	109	9.2 (100)
0653	167	6.6 (118)	45.5 (7)	142	33.1 (103)	48.6 (45)	18	* *	26	* *	148	5.4 (43)
0654	159	5.7 (105)	47.8 (10)	129	17.1 (20)	52.7 (82)	17	* *	24	* *	143	4.2 (16)
0655	149	5.4 (97)	49.0 (16)	132	21.2 (60)	53.0 (86)	11	* *	48	* *	134	14.9 (114)
0657	28	* *	* *	25	* *	* *	4	* *	7	* *	25	* *
0701	151	3.3 (28)	41.7 (3)	137	24.1 (76)	43.1 (21)	12	* *	29	* *	113	4.4 (20)
0702	26	* *	* *	23	* *	* *	3	* *	1	* *	22	* *
0703	100	4.0 (50)	55.0 (75)	89	* *	* *	9	* *	17	* *	65	* *
0704	26	* *	* *	26	* *	* *	3	* *	5	* *	20	* *
0705	52	* *	* *	46	* *	* *	6	* *	9	* *	38	* *
0850	466	3.2 (26)	54.1 (62)	387	11.6 (1)	59.4 (114)	52	* *	71	* *	430	4.4 (20)
0851	347	4.6 (71)	48.1 (12)	296	13.5 (3)	55.4 (102)	45	* *	41	* *	322	5.3 (40)
0901	671	5.5 (100)	54.8 (72)	590	23.9 (74)	49.7 (54)	83	* *	188	23.4 (10)	558	8.6 (96)
0902	27	* *	* *	25	* *	* *	4	* *	11	* *	23	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Maternal Health Indicators by Clinic

Page: 26
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Health Indicators (2, 3)											
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Gestational Diabetes During Pregnancy	Hypertension During Pregnancy	
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester	Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1051	562	3.9 (48)	53.6 (57)	471	23.1 (71)	51.4 (70)	51	* *	131	19.8 (3)	424	5.0 (34)
1101	759	4.6 (71)	55.7 (81)	658	18.1 (32)	53.8 (96)	126	56.3 (36)	212	44.8 (40)	724	6.2 (62)
1102	356	6.2 (115)	50.6 (32)	311	15.8 (9)	57.2 (110)	48	* *	89	* *	327	4.9 (32)
1106	43	* *	* *	33	* *	* *	11	* *	13	* *	38	* *
1250	438	4.8 (77)	52.3 (47)	385	24.9 (82)	47.5 (37)	47	* *	53	* *	397	7.3 (79)
1251	339	4.7 (76)	58.7 (108)	277	19.9 (49)	51.3 (69)	42	* *	61	* *	301	4.0 (11)
1252	211	2.8 (18)	56.9 (95)	148	20.3 (55)	49.3 (49)	26	* *	27	* *	223	13.5 (113)
1253	325	4.0 (50)	51.7 (40)	256	21.1 (59)	48.8 (48)	53	* *	70	* *	329	6.4 (65)
1350	973	3.8 (45)	55.3 (77)	831	20.6 (56)	50.9 (63)	165	34.5 (14)	119	24.4 (14)	882	4.6 (26)
1351	69	* *	* *	60	* *	* *	11	* *	7	* *	62	* *
1352	162	1.2 (3)	58.6 (107)	146	21.2 (60)	54.1 (98)	17	* *	10	* *	147	6.1 (59)
1401	90	* *	* *	71	* *	* *	5	* *	12	* *	60	* *
1402	67	* *	* *	63	* *	* *	5	* *	14	* *	58	* *
1403	9	* *	* *	6	* *	* *	0	* *	3	* *	4	* *
1404	11	* *	* *	5	* *	* *	1	* *	1	* *	5	* *
1501	358	2.0 (10)	58.7 (108)	237	48.1 (113)	35.0 (4)	48	* *	39	* *	393	9.4 (101)
1503	431	6.7 (119)	60.3 (115)	266	44.4 (111)	32.7 (3)	48	* *	71	* *	472	5.7 (49)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Maternal Health Indicators by Clinic

Page: 27
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Health Indicators (2, 3)												
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Gestational Diabetes During Pregnancy	Hypertension During Pregnancy		
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester	Postpartum							
Clinic	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
1504	826	3.0 (20)	53.9 (61)	537	22.7 (68)	48.0 (40)	143	33.6 (12)	183	26.2 (19)	931	8.4 (93)	
1505	362	3.6 (36)	50.0 (25)	319	18.8 (44)	44.5 (26)	48	* *	79	* *	330	2.7 (1)	
1506	216	4.6 (71)	45.4 (6)	153	42.5 (109)	38.6 (8)	28	* *	34	* *	238	8.4 (93)	
1507	236	3.8 (45)	50.0 (25)	190	18.4 (36)	55.8 (103)	29	* *	47	* *	243	7.0 (75)	
1508	38	* *	* *	27	* *	* *	9	* *	5	* *	38	* *	
1509	268	3.4 (31)	56.0 (85)	138	37.7 (105)	38.4 (7)	22	* *	27	* *	257	10.5 (106)	
1510	43	* *	* *	23	* *	* *	2	* *	6	* *	40	* *	
1511	270	3.7 (40)	55.9 (83)	217	23.0 (69)	53.9 (97)	36	* *	41	* *	247	6.9 (74)	
1512	99	* *	* *	65	* *	* *	14	* *	28	* *	97	* *	
1701	301	5.3 (93)	50.5 (31)	243	27.2 (91)	51.0 (65)	15	* *	54	* *	190	5.3 (40)	
1712	19	* *	* *	17	* *	* *	1	* *	1	* *	14	* *	
1950	312	5.8 (108)	52.9 (56)	263	18.3 (35)	53.6 (94)	39	* *	95	* *	279	3.9 (10)	
1952	53	* *	* *	48	* *	* *	7	* *	22	* *	51	* *	
1953	124	4.8 (77)	54.8 (72)	112	21.4 (62)	52.7 (82)	13	* *	48	* *	113	11.5 (109)	
1954	245	4.9 (82)	50.2 (29)	191	29.3 (98)	42.4 (20)	34	* *	103	22.3 (6)	270	5.2 (37)	
1955	52	* *	* *	39	* *	* *	6	* *	24	* *	55	* *	
1957	206	4.9 (82)	55.3 (77)	177	18.6 (39)	50.8 (61)	29	* *	63	* *	186	5.9 (55)	
												186	10.2 (58)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Maternal Health Indicators by Clinic

Page: 28
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Health Indicators (2, 3)											
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Gestational Diabetes During Pregnancy	Hypertension During Pregnancy	
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester	Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1959	63	* *	* *	49	* *	* *	11	* *	8	* *	58	* *
2150	288	1.7 (5)	52.1 (44)	269	36.1 (104)	40.9 (16)	27	* *	72	* *	203	12.3 (111)
2151	121	8.3 (123)	41.3 (2)	105	27.6 (93)	47.6 (38)	10	* *	22	* *	77	* *
2251	188	4.3 (62)	46.8 (8)	166	18.7 (43)	49.4 (51)	9	* *	24	* *	119	3.4 (7)
2252	108	1.9 (7)	51.9 (42)	86	* *	* *	4	* *	12	* *	61	* *
2551	1,293	4.8 (77)	52.4 (48)	428	27.3 (92)	46.0 (33)	274	56.2 (35)	134	38.1 (36)	469	5.3 (40)
2554	38	* *	* *	31	* *	* *	2	* *	3	* *	26	* *
2555	1,554	5.2 (89)	54.2 (64)	728	21.4 (62)	50.0 (56)	336	39.9 (27)	304	24.0 (12)	790	9.1 (99)
3001	394	2.5 (15)	57.6 (98)	332	27.7 (94)	41.6 (17)	50	* *	236	28.8 (26)	299	3.0 (2)
3002	95	* *	* *	69	* *	* *	8	* *	57	* *	72	* *
3003	4	* *	* *	4	* *	* *	1	* *	1	* *	4	* *
3004	67	* *	* *	53	* *	* *	1	* *	12	* *	44	* *
3007	10	* *	* *	8	* *	* *	0	* *	4	* *	7	* *
3009	5	* *	* *	3	* *	* *	0	* *	1	* *	3	* *
3010	2	* *	* *	0	* *	* *	0	* *	0	* *	0	* *
3201	227	4.0 (50)	52.0 (43)	202	24.8 (80)	53.0 (86)	23	* *	34	* *	177	9.6 (102)
3301	1,692	4.4 (66)	54.4 (66)	1,398	17.4 (25)	54.4 (100)	303	28.7 (3)	291	26.8 (20)	1,591	5.2 (37)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Maternal Health Indicators by Clinic

Page: 29
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Health Indicators (2, 3)												
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Gestational Diabetes During Pregnancy	Hypertension During Pregnancy		
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester	Postpartum							
Clinic	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
3302	107	0.0 (1)	60.7 (117)	92	* * * *	18	*	11	* *	101	4.0 (11)	101 11.9(80)	
3304	40	* * * *	* *	36	* * * *	2	*	10	* *	36	* *	36 * *	
3305	39	* * * *	* *	36	* * * *	4	*	10	* *	39	* * 39	* *	
3306	84	* * * *	* *	79	* * * *	11	*	21	* *	80	* * 80	* *	
3308	19	* * * *	* *	13	* * * *	5	*	7	* *	16	* * 16	* *	
3309	12	* * * *	* *	12	* * * *	2	*	3	* *	12	* * 12	* *	
3550	34	* * * *	* *	32	* * * *	5	*	6	* *	23	* * 23	* *	
3551	8	* * * *	* *	8	* * * *	0	*	4	* *	6	* * 6	* *	
3552	13	* * * *	* *	13	* * * *	0	*	4	* *	10	* * 10	* *	
3701	425	5.2 (89)	53.6 (57)	335	17.0 (17)	57.6(112)	65	*	52	* *	393	5.9 (55)	393 7.1(19)
3805	1,351	4.3 (62)	49.5 (20)	1,203	18.0 (30)	53.0 (86)	160	32.5 (10)	134	26.1 (18)	1,246	4.2 (16)	1,244 8.7(44)
3950	842	5.0 (87)	56.7 (93)	705	16.7 (15)	53.6 (94)	119	39.5 (24)	143	35.0 (35)	761	7.5 (83)	760 6.7(14)
3952	5	* * * *	* *	3	* * * *	0	*	0	* *	3	* * 3	* *	
3953	3	* * * *	* *	2	* * * *	0	*	0	* *	2	* * 2	* *	
3954	23	* * * *	* *	21	* * * *	1	*	4	* *	22	* * 22	* *	
4001	429	4.9 (82)	58.3 (104)	346	40.2 (107)	39.0 (10)	56	*	55	* *	408	4.4 (20)	408 11.8(79)
4002	371	4.3 (62)	52.8 (53)	329	26.7 (86)	49.5 (53)	46	*	51	* *	355	10.1(104)	355 5.6(8)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight pre pregnancy (ideal weight gain = 25 to 35 pounds), overweight pre pregnancy (ideal weight gain = 15 to 25 pounds), obese pre pregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Maternal Health Indicators by Clinic

Page: 30
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Health Indicators (2, 3)											
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Gestational Diabetes During Pregnancy	Hypertension During Pregnancy	
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester	Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4103	32	* *	* *	27	* *	* *	4	* *	4	* *	32	* *
4106	307	2.6 (16)	60.9 (119)	250	22.4 (67)	48.0 (40)	48	* *	57	* *	306	6.5 (70)
4107	1,449	3.7 (40)	50.2 (29)	1,132	19.5 (46)	51.9 (77)	218	45.0 (31)	260	25.8 (17)	1,384	5.2 (37)
4109	400	2.3 (12)	54.3 (65)	208	47.1 (112)	36.5 (5)	66	* *	93	* *	459	5.7 (49)
4110	636	3.0 (20)	55.0 (75)	462	24.0 (75)	43.3 (24)	81	* *	93	* *	656	4.0 (11)
4112	238	1.7 (5)	64.7 (123)	217	69.1 (115)	12.0 (1)	18	* *	14	* *	237	11.8 (110)
4113	255	1.6 (4)	57.3 (96)	227	26.4 (85)	37.4 (6)	28	* *	20	* *	245	5.7 (49)
4120	777	4.8 (77)	50.6 (32)	524	20.2 (53)	49.8 (55)	124	34.7 (16)	177	24.9 (15)	780	4.1 (15)
4160	587	5.3 (93)	52.6 (52)	315	29.8 (101)	42.2 (19)	141	31.2 (9)	131	22.9 (8)	683	6.4 (65)
4351	242	3.3 (28)	52.5 (50)	219	19.6 (47)	53.4 (92)	21	* *	60	* *	181	4.4 (20)
4352	71	* *	* *	63	* *	* *	8	* *	20	* *	52	* *
4353	190	4.2 (59)	56.8 (94)	164	39.0 (106)	40.2 (14)	14	* *	33	* *	145	7.6 (85)
4354	335	4.2 (59)	52.8 (53)	294	24.1 (76)	52.4 (79)	31	* *	68	* *	241	5.4 (43)
4355	144	6.3 (116)	52.1 (44)	122	13.1 (2)	62.3 (115)	12	* *	36	* *	97	* *
4356	206	5.8 (108)	59.2 (111)	182	21.4 (62)	51.1 (66)	17	* *	28	* *	157	5.7 (49)
4357	3	* *	* *	3	* *	* *	0	* *	0	* *	2	* *
4358	83	* *	* *	64	* *	* *	9	* *	15	* *	63	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Maternal Health Indicators by Clinic

Page: 31
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Health Indicators (2, 3)											
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Gestational Diabetes During Pregnancy	Hypertension During Pregnancy	
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester	Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4359	221	6.8 (120)	58.8 (110)	193	23.3 (72)	48.7 (47)	17	* *	43	* *	158	7.0 (75)
4360	63	* *	* *	58	* *	* *	5	* *	19	* *	44	* *
4361	109	5.5 (100)	54.1 (62)	96	* *	* *	14	* *	22	* *	75	* *
4362	143	3.5 (34)	49.0 (16)	126	15.9 (10)	58.7 (113)	14	* *	42	* *	104	4.8 (30)
4363	357	3.6 (36)	49.0 (16)	321	17.4 (25)	55.8 (103)	26	* *	94	* *	261	6.5 (70)
4401	559	6.4 (117)	51.2 (36)	492	19.7 (48)	49.4 (51)	45	* *	132	22.0 (5)	407	5.9 (55)
4501	378	3.4 (31)	56.6 (90)	348	24.7 (79)	46.8 (36)	32	* *	64	* *	308	7.8 (87)
4601	593	4.0 (50)	56.0 (85)	533	28.1 (95)	40.0 (13)	106	20.8 (1)	132	29.5 (29)	587	8.2 (91)
4701	525	3.8 (45)	50.1 (28)	477	17.2 (22)	56.6 (107)	90	* *	182	18.1 (2)	473	6.1 (59)
5050	564	5.7 (105)	51.6 (37)	177	18.6 (39)	54.2 (99)	132	36.4 (20)	82	* *	166	10.2 (105)
5051	1,086	4.8 (77)	53.8 (59)	408	14.2 (4)	56.6 (107)	219	34.7 (16)	151	27.2 (21)	390	6.4 (65)
5052	886	4.0 (50)	54.4 (66)	269	18.6 (39)	53.5 (93)	184	39.7 (25)	112	24.1 (13)	257	7.8 (87)
5101	918	3.5 (34)	49.6 (21)	763	16.0 (11)	53.3 (91)	183	39.3 (23)	578	29.1 (27)	794	7.4 (81)
5202	258	1.9 (7)	51.6 (37)	232	15.1 (5)	50.0 (56)	36	* *	65	* *	183	8.2 (91)
5203	66	* *	* *	58	* *	* *	3	* *	15	* *	41	* *
5204	73	* *	* *	62	* *	* *	2	* *	20	* *	46	* *
5601	476	3.4 (31)	54.4 (66)	433	16.6 (14)	56.1 (106)	41	* *	101	34.7 (33)	377	7.2 (78)
											376	12.5 (87)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Maternal Health Indicators by Clinic

Page: 32
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Health Indicators (2, 3)											
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Gestational Diabetes During Pregnancy	Hypertension During Pregnancy	
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester	Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
5801	922	5.3 (93)	51.8 (41)	784	23.6 (73)	50.9 (63)	115	28.7 (3)	224	23.7 (11)	828	6.6 (72)
5803	17	* *	* *	16	* *	* *	0	* *	9	* *	15	* *
5804	6	* *	* *	6	* *	* *	0	* *	2	* *	7	* *
6101	17	* *	* *	15	* *	* *	4	* *	0	* *	15	* *
6102	59	* *	* *	54	* *	* *	6	* *	7	* *	63	* *
6103	1,568	4.0 (50)	55.7 (81)	1,359	16.3 (12)	52.5 (80)	204	38.2 (22)	104	30.8 (30)	1,524	6.4 (65)
6104	35	* *	* *	25	* *	* *	6	* *	5	* *	36	* *
6105	16	* *	* *	16	* *	* *	2	* *	1	* *	18	* *
6106	1	* *	* *	1	* *	* *	0	* *	0	* *	3	* *
6301	1,942	4.5 (70)	52.2 (46)	1,716	17.0 (17)	50.8 (61)	354	35.3 (18)	701	23.1 (9)	1,660	6.3 (64)
6312	1,041	3.6 (36)	57.6 (98)	937	17.1 (20)	51.4 (70)	209	33.0 (11)	378	29.1 (27)	903	7.5 (83)
6316	590	5.6 (104)	52.5 (50)	539	16.5 (13)	52.5 (80)	104	27.9 (2)	238	20.2 (4)	503	4.6 (26)
6320	96	* *	* *	79	* *	* *	16	* *	28	* *	73	* *
6325	59	* *	* *	47	* *	* *	10	* *	19	* *	45	* *
6330	205	2.4 (14)	54.6 (71)	153	20.9 (58)	46.4 (34)	33	* *	44	* *	167	6.0 (58)
6335	242	5.4 (97)	48.8 (14)	206	17.5 (27)	48.1 (42)	34	* *	53	* *	206	4.4 (20)
6340	56	* *	* *	41	* *	* *	5	* *	16	* *	48	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Maternal Health Indicators by Clinic

Page: 33
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Health Indicators (2, 3)												
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Gestational Diabetes During Pregnancy	Hypertension During Pregnancy		
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester	Postpartum							
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
6345	21	* *	* *	17	* *	* *	4	* *	8	* *	18	* *	
6350	110	5.5 (100)	50.0 (25)	89	* *	* *	17	* *	26	* *	91	* *	
7304	74	* *	* *	54	* *	* *	9	* *	15	* *	70	* *	
7306	945	4.6 (71)	55.9 (83)	750	17.3 (23)	50.5 (59)	138	37.0 (21)	136	27.9 (23)	785	7.1 (77)	
7308	73	* *	* *	69	* *	* *	9	* *	12	* *	66	* *	
7310	23	* *	* *	21	* *	* *	4	* *	1	* *	19	* *	
7401	862	4.9 (82)	49.7 (22)	760	18.2 (34)	52.9 (85)	137	30.7 (8)	467	22.5 (7)	718	11.1 (108)	
7402	53	* *	* *	51	* *	* *	12	* *	22	* *	47	* *	
7403	26	* *	* *	25	* *	* *	5	* *	15	* *	21	* *	
7404	25	* *	* *	22	* *	* *	0	* *	19	* *	16	* *	
7601	290	5.9 (112)	52.8 (53)	243	15.6 (7)	52.7 (82)	28	* *	69	* *	229	10.0 (103)	
7850	378	5.8 (108)	51.6 (37)	331	17.5 (27)	55.0 (101)	38	* *	64	* *	347	4.3 (19)	
7851	54	* *	* *	50	* *	* *	11	* *	6	* *	50	* *	
7852	39	* *	* *	38	* *	* *	3	* *	9	* *	38	* *	
8150	19	* *	* *	15	* *	* *	0	* *	4	* *	18	* *	
8151	1,177	3.7 (40)	54.8 (72)	951	25.9 (84)	44.5 (26)	213	40.8 (28)	305	27.5 (22)	1,050	8.1 (90)	
8152	180	7.2 (122)	43.3 (5)	126	28.6 (96)	57.1 (109)	25	* *	43	* *	184	3.3 (6)	
												184	19.0 (115)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Maternal Health Indicators by Clinic

Page: 34
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Health Indicators (2, 3)											
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Gestational Diabetes During Pregnancy	Hypertension During Pregnancy	
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester	Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8153	9	* *	* *	7	* *	* *	3	* *	6	* *	8	* *
8154	103	6.8 (120)	33.0 (1)	86	* *	* *	12	* *	35	* *	87	* *
8155	20	* *	* *	18	* *	* *	3	* *	6	* *	19	* *
8156	9	* *	* *	8	* *	* *	3	* *	2	* *	7	* *
8157	13	* *	* *	7	* *	* *	1	* *	4	* *	10	* *
8201	1,000	5.0 (87)	52.4 (48)	604	16.7 (15)	55.8 (103)	255	29.4 (7)	237	24.9 (15)	588	7.8 (87)
8202	1,131	4.0 (50)	46.9 (9)	677	26.7 (86)	39.0 (10)	181	35.4 (19)	223	28.7 (25)	839	5.4 (43)
8203	132	2.3 (12)	56.1 (87)	54	* *	* *	21	* *	27	* *	52	* *
8250	324	4.6 (71)	49.4 (19)	201	15.4 (6)	50.7 (60)	98	* *	68	* *	213	6.1 (59)
8255	960	5.5 (100)	51.0 (35)	253	28.9 (97)	45.8 (31)	249	28.9 (5)	172	16.9 (1)	312	6.4 (65)
8257	208	5.8 (108)	48.6 (13)	103	30.1 (102)	38.8 (9)	40	* *	40	* *	134	6.7 (73)
8258	590	3.7 (40)	55.4 (79)	344	29.4 (99)	44.8 (28)	130	29.2 (6)	91	* *	326	7.4 (81)
8260	180	4.4 (66)	42.8 (4)	157	24.8 (80)	40.8 (15)	42	* *	41	* *	158	12.7 (112)
8901	343	3.2 (26)	60.9 (119)	257	26.8 (88)	48.6 (45)	55	* *	32	* *	292	3.1 (3)
8902	219	3.7 (40)	57.5 (97)	176	19.3 (45)	51.1 (66)	33	* *	36	* *	206	8.7 (98)
8903	439	3.9 (48)	60.1 (113)	343	20.1 (51)	49.3 (49)	76	* *	44	* *	387	5.7 (49)
8904	315	1.9 (7)	63.8 (122)	268	26.9 (89)	44.0 (25)	49	* *	51	* *	276	10.9 (107)
											277	14.4 (110)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight pre pregnancy (ideal weight gain = 25 to 35 pounds), overweight pre pregnancy (ideal weight gain = 15 to 25 pounds), obese pre pregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Maternal Health Indicators by Clinic

Page: 35
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Health Indicators (2, 3)											
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Gestational Diabetes During Pregnancy	Hypertension During Pregnancy	
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester	Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8905	1,001	3.6 (36)	56.6 (90)	822	25.5 (83)	43.1 (21)	87	* *	93	* *	887	4.2 (16)
9004	4	* * * *	* *	3	* * * *	* *	2	* *	0	* *	3	* * *
9012	13	* * * *	* *	10	* * * *	* *	0	* *	0	* *	13	* * 13
9013	27	* * * *	* *	23	* * * *	* *	4	* *	1	* *	30	* * 30
9015	8	* * * *	* *	7	* * * *	* *	2	* *	0	* *	10	* * 10
9016	33	* * * *	* *	25	* * * *	* *	4	* *	5	* *	30	* * 30
9017	8	* * * *	* *	6	* * * *	* *	1	* *	1	* *	9	* * 9
9018	30	* * * *	* *	28	* * * *	* *	0	* *	1	* *	31	* * 31
9019	12	* * * *	* *	10	* * * *	* *	1	* *	0	* *	12	* * 12
9021	14	* * * *	* *	13	* * * *	* *	3	* *	0	* *	20	* * 20
9040	7	* * * *	* *	5	* * * *	* *	0	* *	1	* *	6	* * 6
9041	155	5.2 (89)	58.1 (102)	135	17.0 (17)	51.1 (66)	8	* *	7	* *	141	5.7 (49)
9045	474	5.7 (105)	53.8 (59)	400	18.5 (37)	51.5 (75)	65	* *	63	* *	427	3.5 (8)
9050	775	2.7 (17)	59.7 (112)	685	22.2 (66)	45.8 (31)	110	34.5 (14)	147	28.6 (24)	696	5.6 (48)
9051	912	2.2 (11)	56.6 (90)	721	21.8 (65)	46.7 (35)	66	* *	134	32.1 (32)	807	7.6 (85)
9053	1,087	4.2 (59)	60.6 (116)	963	15.6 (7)	57.2 (110)	206	39.8 (26)	358	31.6 (31)	967	4.0 (11)
9055	43	* * * *	* *	40	* * * *	* *	4	* *	0	* *	21	* * 22

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Maternal Health Indicators by Clinic

Page: 36
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Health Indicators (2, 3)											
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Gestational Diabetes During Pregnancy	Hypertension During Pregnancy	
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester	Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
9058	252	4.4 (66)	48.0 (11)	227	49.3 (114)	22.9 (2)	31	* *	48	* *	226	15.5 (115)
9066	919	4.4 (66)	57.6 (98)	823	20.7 (57)	48.5 (44)	156	41.7 (29)	152	46.7 (41)	840	5.0 (34)
9068	562	4.3 (62)	56.2 (88)	524	24.2 (78)	48.3 (43)	88	* *	111	56.8 (43)	518	5.0 (34)
9069	1,066	3.1 (24)	60.8 (118)	946	18.1 (32)	51.4 (70)	199	34.2 (13)	310	34.8 (34)	997	4.8 (30)
9070	802	6.1 (114)	55.5 (80)	695	19.9 (49)	50.1 (58)	105	53.3 (33)	145	40.0 (39)	741	4.9 (32)
9072	714	3.1 (24)	60.1 (113)	669	18.5 (37)	53.1 (89)	148	41.9 (30)	216	47.7 (42)	660	5.5 (46)
9074	689	3.0 (20)	61.0 (121)	603	17.6 (29)	53.2 (90)	151	55.0 (34)	212	39.2 (37)	620	4.5 (25)
9080	869	4.0 (50)	58.0 (101)	739	17.3 (23)	51.4 (70)	144	52.8 (32)	177	39.5 (38)	786	6.2 (62)
Unknown Clinics	21	* * * *	*	14	* * * *	*	1	* *	7	* *	15	* *
Michigan	64,771	4.2	54.2	50,312	21.8	49.4	9,832	36.8	13,734	27.6	53,787	6.3
Nation (Prior Year)	1,224,766	4.7	52.1	963,624	21.4	48.2	188,530	33.8	416,710	29.6	346,704	4.4

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Maternal Behavioral Indicators by Clinic

Page: 37
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Behavioral Indicators (2, 3)									
	Medical Care			WIC Enrollment			Multivitamin Consumption			
	1st Trimester	No Care	1st Trimester	Prepregnancy (4)		During Pregnancy (5)				
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
0101	51	*	*	60	*	*	9	*	*	11
0102	162	93.2 (13)	0.6 (33)	173	43.9 (26)	38	*	*	45	*
0103	172	93.6 (11)	1.7 (97)	176	50.6 (6)	38	*	*	38	*
0104	50	*	*	52	*	*	12	*	*	13
0250	56	*	*	59	*	*	9	*	*	11
0251	52	*	*	62	*	*	12	*	*	15
0252	44	*	*	48	*	*	6	*	*	9
0253	65	*	*	65	*	*	13	*	*	16
0401	214	89.7 (53)	0.9 (59)	231	45.9 (17)	61	*	*	64	*
0402	161	95.7 (2)	0.6 (33)	166	42.8 (32)	35	*	*	38	*
0403	51	*	*	55	*	*	6	*	*	6
0404	54	*	*	53	*	*	9	*	*	11
0501	199	92.0 (24)	2.0 (104)	216	55.6 (2)	43	*	*	48	*
0502	1	*	*	1	*	*	0	*	*	0
0503	75	*	*	78	*	*	17	*	*	17
0504	212	96.7 (1)	0.5 (24)	224	51.8 (5)	43	*	*	46	*
0505	135	92.6 (19)	3.0 (112)	143	53.8 (4)	38	*	*	40	*
0506	65	*	*	74	*	*	21	*	*	22
0650	446	92.4 (21)	0.7 (46)	472	40.7 (47)	232	21.6 (18)	238	89.1 (26)	
0651	244	86.9 (75)	2.9 (110)	263	41.8 (41)	135	20.7 (24)	151	79.5 (73)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Maternal Behavioral Indicators by Clinic

Page: 38
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Behavioral Indicators (2, 3)									
	Medical Care			WIC Enrollment			Multivitamin Consumption			
	1st Trimester	No Care	1st Trimester	Prepregnancy (4)		During Pregnancy (5)				
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
0652	113	94.7 (5)	0.9 (59)	122	39.3 (57)	57	*	67	*	*
0653	159	91.8 (25)	1.9 (102)	170	42.9 (31)	97	*	99	*	*
0654	158	87.3 (71)	3.2 (113)	165	55.2 (3)	90	*	93	*	*
0655	140	90.7 (41)	1.4 (91)	148	35.1 (75)	73	*	76	*	*
0657	27	*	*	28	*	17	*	17	*	*
0701	149	84.6 (89)	0.0 (1)	172	41.9 (39)	27	*	27	*	*
0702	26	*	*	30	*	11	*	11	*	*
0703	90	*	*	99	*	20	*	22	*	*
0704	24	*	*	25	*	5	*	5	*	*
0705	56	*	*	60	*	15	*	15	*	*
0850	444	91.2 (32)	0.0 (1)	481	39.7 (52)	356	16.0 (47)	360	84.2 (54)	
0851	333	92.8 (16)	0.6 (33)	356	39.6 (54)	275	13.5 (58)	282	83.7 (55)	
0901	667	91.2 (32)	1.0 (68)	714	41.2 (46)	158	22.2 (15)	244	91.4 (16)	
0902	27	*	*	28	*	6	*	6	*	*
1051	506	89.5 (57)	2.6 (109)	582	45.7 (18)	131	25.2 (7)	143	85.3 (46)	
1101	721	79.2 (100)	1.2 (80)	821	34.8 (77)	485	10.7 (65)	489	89.4 (25)	
1102	332	90.7 (41)	0.0 (1)	367	37.6 (69)	193	15.0 (54)	210	94.8 (2)	
1106	37	*	*	45	*	26	*	26	*	*
1250	408	90.4 (47)	0.5 (24)	439	49.2 (9)	238	23.9 (10)	287	68.6 (81)	
1251	317	90.5 (45)	0.9 (59)	346	42.8 (32)	214	16.4 (43)	230	81.7 (65)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Maternal Behavioral Indicators by Clinic

Page: 39
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Behavioral Indicators (2, 3)									
	Medical Care			WIC Enrollment			Multivitamin Consumption			
	1st Trimester	No Care	1st Trimester	Prepregnancy (4)		During Pregnancy (5)				
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
1252	227	82.4 (96)	5.3 (116)	250	45.6 (19)	194	22.7 (13)	197	74.1 (80)	
1253	343	86.9 (75)	3.2 (113)	374	35.0 (76)	212	22.2 (15)	217	82.9 (60)	
1350	903	92.1 (23)	1.8 (100)	1,010	37.9 (64)	760	31.1 (4)	791	82.6 (63)	
1351	65	*	*	71	*	57	*	58	*	*
1352	150	90.0 (52)	0.7 (46)	165	50.3 (8)	139	17.3 (36)	139	81.3 (67)	
1401	82	*	*	93	*	18	*	18	*	*
1402	64	*	*	71	*	17	*	17	*	*
1403	8	*	*	9	*	2	*	2	*	*
1404	7	*	*	12	*	3	*	3	*	*
1501	398	85.9 (82)	1.3 (85)	471	38.2 (63)	256	18.0 (34)	278	74.8 (79)	
1503	406	83.0 (94)	1.0 (68)	567	37.4 (71)	197	33.5 (2)	338	79.9 (69)	
1504	952	92.6 (19)	1.1 (74)	1,004	39.1 (58)	637	15.4 (50)	669	82.7 (62)	
1505	337	95.3 (3)	0.6 (33)	362	37.8 (66)	237	28.3 (6)	239	92.1 (11)	
1506	243	90.5 (45)	2.9 (110)	265	42.3 (37)	171	9.9 (69)	173	79.8 (70)	
1507	251	94.0 (8)	1.2 (80)	264	39.8 (51)	164	21.3 (20)	169	85.2 (48)	
1508	43	*	*	71	*	28	*	31	*	*
1509	261	85.8 (84)	2.3 (108)	305	47.9 (13)	156	18.6 (33)	170	76.5 (78)	
1510	42	*	*	46	*	34	*	34	*	*
1511	256	93.0 (14)	0.4 (22)	278	40.3 (48)	182	18.7 (32)	182	85.2 (48)	
1512	98	*	*	107	32.7 (78)	66	*	66	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Maternal Behavioral Indicators by Clinic

Page: 40
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Behavioral Indicators (2, 3)									
	Medical Care			WIC Enrollment			Multivitamin Consumption			
	1st Trimester	No Care	1st Trimester	Prepregnancy (4)		During Pregnancy (5)				
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
1701	267	86.1 (79)	1.1 (74)	305	48.2 (12)	26	*	40	*	*
1712	17	*	*	21	*	3	*	4	*	*
1950	289	94.8 (4)	0.0 (1)	317	40.1 (49)	168	19.6 (28)	172	93.0 (8)	
1952	50	*	*	53	*	20	*	22	*	*
1953	116	91.4 (31)	0.9 (59)	122	42.6 (34)	59	*	59	*	*
1954	286	90.9 (37)	0.7 (46)	290	41.7 (42)	151	17.2 (38)	169	82.8 (61)	
1955	56	*	*	59	*	29	*	32	*	*
1957	194	91.8 (25)	0.5 (24)	212	39.6 (54)	99	*	101	85.1 (50)	
1959	59	*	*	66	*	33	*	35	*	*
2150	275	93.8 (10)	0.7 (46)	290	41.7 (42)	47	*	51	*	*
2151	104	86.5 (78)	1.9 (102)	125	42.4 (35)	22	*	28	*	*
2251	176	91.5 (29)	0.6 (33)	184	50.5 (7)	24	*	34	*	*
2252	98	*	*	109	64.2 (1)	14	*	20	*	*
2551	530	80.0 (99)	0.8 (54)	1,347	27.5 (88)	379	6.1 (77)	494	85.0 (51)	
2554	31	*	*	38	*	12	*	12	*	*
2555	891	88.1 (67)	0.0 (1)	1,653	24.9 (97)	484	11.6 (63)	591	85.8 (45)	
3001	366	77.0 (107)	1.1 (74)	392	47.4 (14)	111	6.3 (75)	113	83.2 (59)	
3002	95	*	*	98	*	28	*	28	*	*
3003	4	*	*	4	*	1	*	1	*	*
3004	51	*	*	70	*	23	*	23	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Maternal Behavioral Indicators by Clinic

Page: 41
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Behavioral Indicators (2, 3)									
	Medical Care			WIC Enrollment			Multivitamin Consumption			
	1st Trimester	No Care	1st Trimester	Prepregnancy (4)		During Pregnancy (5)				
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
3007	8	*	*	11	*	*	2	*	*	2
3009	3	*	*	9	*	*	0	*	*	0
3010	0	*	*	3	*	*	0	*	*	0
3201	223	92.8 (16)	0.0 (1)	230	38.3 (60)	71	*	*	73	*
3301	1,609	86.7 (77)	1.2 (80)	1,788	27.0 (90)	1,263	16.8 (40)	1,431	86.4 (41)	
3302	98	*	*	109	23.9 (100)	45	*	*	86	*
3304	37	*	*	39	*	*	29	*	*	30
3305	38	*	*	40	*	*	25	*	*	26
3306	80	*	*	87	*	*	54	*	*	57
3308	18	*	*	19	*	*	12	*	*	13
3309	12	*	*	12	*	*	8	*	*	8
3550	32	*	*	32	*	*	9	*	*	10
3551	9	*	*	9	*	*	1	*	*	1
3552	14	*	*	14	*	*	0	*	*	0
3701	405	92.8 (16)	0.2 (16)	438	36.5 (72)	155	12.3 (62)	217	91.7 (14)	
3805	1,257	89.7 (53)	0.8 (54)	1,354	43.6 (27)	1,026	15.8 (48)	1,114	76.9 (77)	
3950	769	89.7 (53)	0.9 (59)	857	37.7 (67)	497	12.7 (60)	502	85.3 (46)	
3952	4	*	*	5	*	*	2	*	*	2
3953	2	*	*	3	*	*	1	*	*	1
3954	24	*	*	24	*	*	15	*	*	15

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Maternal Behavioral Indicators by Clinic

Page: 42
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Behavioral Indicators (2, 3)							
	Medical Care			WIC Enrollment		Multivitamin Consumption		
	1st Trimester	No Care	1st Trimester	Prepregnancy (4)		During Pregnancy (5)		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number
4001	418	83.0 (94)	1.7 (97)	467	48.8 (11)	263	12.5 (61)	267
4002	363	87.6 (68)	0.6 (33)	383	41.3 (44)	218	21.1 (23)	225
4103	30	*	*	32	*	15	*	15
4106	307	87.0 (73)	1.0 (68)	331	29.0 (86)	157	8.9 (71)	172
4107	1,421	88.6 (63)	1.3 (85)	1,534	36.3 (73)	812	21.6 (18)	974
4109	457	90.4 (47)	1.8 (100)	506	35.8 (74)	275	22.9 (11)	281
4110	658	89.1 (59)	2.1 (107)	715	38.3 (60)	366	15.3 (52)	451
4112	239	62.8 (116)	0.0 (1)	250	37.6 (69)	150	24.7 (8)	174
4113	215	78.6 (103)	3.3 (115)	266	47.0 (15)	141	2.8 (80)	159
4120	801	89.6 (56)	0.9 (59)	849	30.0 (82)	511	16.8 (40)	522
4160	691	88.7 (62)	0.7 (46)	739	26.3 (92)	463	21.2 (22)	479
4351	233	88.8 (60)	1.3 (85)	247	44.5 (23)	65	*	68
4352	66	*	*	73	*	17	*	19
4353	184	87.5 (69)	1.6 (94)	194	43.3 (29)	48	*	53
4354	316	94.0 (8)	1.3 (85)	328	42.4 (35)	85	*	96
4355	124	84.7 (88)	1.6 (94)	155	41.3 (44)	40	*	43
4356	208	94.2 (7)	0.5 (24)	232	44.0 (24)	62	*	63
4357	4	*	*	3	*	0	*	0
4358	71	*	*	91	*	21	*	32
4359	212	85.8 (84)	0.9 (59)	232	44.8 (22)	54	*	54

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Maternal Behavioral Indicators by Clinic

Page: 43
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Behavioral Indicators (2, 3)									
	Medical Care			WIC Enrollment			Multivitamin Consumption			
	1st Trimester	No Care	1st Trimester	Prepregnancy (4)		During Pregnancy (5)				
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
4360	58	*	*	62	*	*	13	*	*	17
4361	98	*	*	109	44.0 (24)	28	*	*	29	*
4362	131	92.4 (21)	0.0 (1)	148	46.6 (16)	28	*	*	29	*
4363	328	90.9 (37)	1.5 (93)	363	41.9 (39)	93	*	*	119	88.2 (32)
4401	509	88.4 (64)	0.8 (54)	556	25.2 (95)	131	22.9 (11)	141	84.4 (53)	
4501	372	91.1 (34)	1.6 (94)	383	43.1 (30)	106	21.7 (17)	109	95.4 (1)	
4601	620	91.1 (34)	0.2 (16)	669	23.6 (101)	244	15.6 (49)	315	89.8 (22)	
4701	494	90.9 (37)	1.2 (80)	533	29.6 (84)	226	19.9 (27)	228	89.0 (28)	
5050	189	91.5 (29)	0.0 (1)	577	23.2 (103)	184	17.9 (35)	242	92.6 (9)	
5051	435	90.1 (51)	0.5 (24)	1,133	30.0 (82)	336	20.5 (25)	460	89.1 (26)	
5052	287	86.1 (79)	0.3 (19)	903	25.5 (94)	304	16.4 (43)	409	90.0 (20)	
5101	855	90.2 (50)	0.2 (16)	959	22.6 (104)	278	24.1 (9)	361	94.2 (6)	
5202	244	94.3 (6)	0.0 (1)	263	45.2 (21)	55	*	58	*	*
5203	62	*	*	70	*	*	11	*	11	*
5204	67	*	*	78	*	*	11	*	11	*
5601	468	93.4 (12)	0.6 (33)	477	39.4 (56)	130	22.3 (14)	135	89.6 (23)	
5801	873	87.2 (72)	1.3 (85)	952	37.7 (67)	454	19.2 (31)	456	91.7 (14)	
5803	17	*	*	20	*	*	1	*	1	*
5804	7	*	*	7	*	*	0	*	0	*
6101	16	*	*	17	*	*	10	*	11	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Maternal Behavioral Indicators by Clinic

Page: 44
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Behavioral Indicators (2, 3)									
	Medical Care			WIC Enrollment			Multivitamin Consumption			
	1st Trimester	No Care	1st Trimester	Prepregnancy (4)		During Pregnancy (5)				
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
6102	62	*	*	64	*	*	41	*	*	41
6103	1,570	90.7 (41)	0.6 (33)	1,704	49.1 (10)	865	15.4 (50)	946	79.6 (72)	
6104	37	*	*	41	*	*	23	*	*	24
6105	18	*	*	18	*	*	13	*	*	13
6106	3	*	*	3	*	*	1	*	*	1
6301	1,794	85.9 (82)	1.0 (68)	1,971	22.0 (105)	644	16.5 (42)	770	84.9 (52)	
6312	975	86.1 (79)	0.9 (59)	1,060	13.3 (119)	347	19.6 (28)	416	90.9 (19)	
6316	553	90.6 (44)	0.7 (46)	593	18.7 (114)	183	21.3 (20)	226	88.1 (33)	
6320	88	*	*	100	21.0 (109)	7	*	35	*	*
6325	51	*	*	59	*	*	15	*	*	21
6330	181	74.0 (114)	1.7 (97)	211	32.7 (78)	74	*	86	*	*
6335	231	90.9 (37)	0.9 (59)	257	30.7 (80)	16	*	99	*	*
6340	53	*	*	58	*	*	4	*	*	16
6345	19	*	*	21	*	*	1	*	*	7
6350	98	*	*	115	20.0 (111)	16	*	44	*	*
7304	79	*	*	92	*	*	20	*	*	22
7306	979	84.9 (86)	1.0 (68)	1,081	40.1 (49)	347	19.6 (28)	361	83.7 (55)	
7308	74	*	*	80	*	*	28	*	*	30
7310	25	*	*	26	*	*	6	*	*	6
7401	769	88.4 (64)	2.0 (104)	907	38.8 (59)	315	15.2 (53)	346	81.2 (68)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Maternal Behavioral Indicators by Clinic

Page: 45
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Behavioral Indicators (2, 3)									
	Medical Care			WIC Enrollment			Multivitamin Consumption			
	1st Trimester	No Care	1st Trimester	Prepregnancy (4)		During Pregnancy (5)				
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
7402	52	*	*	56	*	*	18	*	*	19
7403	25	*	*	28	*	*	4	*	*	5
7404	22	*	*	26	*	*	7	*	*	9
7601	263	91.6 (28)	1.1 (74)	295	39.7 (52)	89	*	*	97	*
7850	356	91.0 (36)	0.0 (1)	386	43.5 (28)	177	16.4 (43)	177	88.1 (33)	
7851	51	*	*	53	*	*	36	*	*	36
7852	37	*	*	39	*	*	16	*	*	16
8150	18	*	*	20	*	*	8	*	*	8
8151	1,112	87.4 (70)	0.6 (33)	1,208	29.4 (85)	559	20.2 (26)	603	87.7 (38)	
8152	185	87.0 (73)	0.0 (1)	198	37.9 (64)	98	*	*	101	86.1 (43)
8153	9	*	*	10	*	*	3	*	*	4
8154	92	*	*	105	25.7 (93)	38	*	*	39	*
8155	19	*	*	22	*	*	4	*	*	8
8156	8	*	*	10	*	*	5	*	*	5
8157	10	*	*	14	*	*	7	*	*	10
8201	646	88.2 (66)	0.3 (19)	1,018	18.8 (113)	465	14.6 (55)	482	92.3 (10)	
8202	889	89.3 (58)	1.0 (68)	1,213	42.2 (38)	318	16.4 (43)	510	81.8 (64)	
8203	59	*	*	137	13.9 (118)	60	*	*	73	*
8250	217	81.1 (97)	0.0 (1)	341	12.0 (120)	243	14.0 (56)	245	94.7 (3)	
8255	349	90.3 (49)	0.6 (33)	985	19.9 (112)	448	17.0 (39)	515	93.8 (7)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Maternal Behavioral Indicators by Clinic

Page: 46
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Behavioral Indicators (2, 3)							
	Medical Care			WIC Enrollment		Multivitamin Consumption		
	1st Trimester	No Care	1st Trimester	Prepregnancy (4)		During Pregnancy (5)		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number
8257	144	91.7 (27)	0.7 (46)	215	25.1 (96)	104	2.9 (79)	111
8258	367	88.8 (60)	0.8 (54)	603	24.0 (99)	243	17.3 (36)	252
8260	168	83.3 (92)	0.0 (1)	182	26.4 (91)	66	* *	84
8901	339	75.5 (112)	0.6 (33)	363	21.8 (106)	110	10.0 (68)	145
8902	241	78.8 (102)	0.4 (22)	255	21.6 (107)	42	* *	96
8903	437	75.7 (111)	0.5 (24)	462	20.8 (110)	118	7.6 (74)	201
8904	329	83.6 (91)	0.6 (33)	345	21.2 (108)	71	* *	141
8905	1,016	84.8 (87)	0.3 (19)	1,088	30.7 (80)	225	6.2 (76)	442
9004	3	*	*	4	*	0	*	3
9012	14	*	*	13	*	9	*	10
9013	30	*	*	29	*	12	*	12
9015	10	*	*	10	*	8	*	9
9016	31	*	*	34	*	21	*	21
9017	9	*	*	10	*	6	*	8
9018	33	*	*	33	*	19	*	19
9019	13	*	*	14	*	6	*	10
9021	20	*	*	20	*	11	*	14
9040	6	*	*	7	*	0	*	3
9041	168	80.4 (98)	0.6 (33)	174	24.1 (98)	30	*	35
9045	449	75.3 (113)	2.0 (104)	492	28.9 (87)	224	8.0 (72)	227
								83.3 (58)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Maternal Behavioral Indicators by Clinic

Page: 47
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Behavioral Indicators (2, 3)					
	Medical Care		WIC Enrollment		Multivitamin Consumption	
	1st Trimester	No Care	1st Trimester		Prepregnancy (4)	During Pregnancy (5)
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number % (Rank)
9050	725	83.2 (93)	0.8 (54)	785	27.1 (89)	210 32.4 (3)
9051	830	83.9 (90)	0.7 (46)	936	45.3 (20)	228 44.3 (1)
9053	1,008	78.1 (105)	0.5 (24)	1,099	14.4 (117)	375 30.7 (5)
9055	49	*	*	52	*	1 *
9058	244	93.0 (14)	0.0 (1)	253	38.3 (60)	127 4.7 (78)
9066	876	75.9 (110)	1.1 (74)	928	11.0 (122)	215 10.2 (66)
9068	542	76.9 (108)	1.3 (85)	554	15.9 (116)	158 10.8 (64)
9069	1,013	79.1 (101)	0.5 (24)	1,083	10.7 (123)	417 12.9 (59)
9070	763	77.5 (106)	0.5 (24)	815	17.8 (115)	225 13.8 (57)
9072	688	73.5 (115)	1.2 (80)	717	10.5 (124)	330 7.9 (73)
9074	646	76.6 (109)	1.1 (74)	687	11.8 (121)	258 10.1 (67)
9080	839	78.4 (104)	1.4 (91)	907	23.3 (102)	381 9.2 (70)
Unknown Clinics	23	*	*	51	*	8 *
Michigan	58,275	86.8	1.0	68,253	32.5	27,982 18.0
Nation (Prior Year)	580,485	80.1	3.1	1,187,387	31.7	266,567 19.3
						269,548 80.6

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Smoking Indicators by Clinic

(1)

Page: 48
Run Date: 7 / 21 / 10

Clinic	Comparison of Smoking Indicators (2, 3)											
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household						
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum				
Clinic	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)		
0101	46	* *	55	* *	13	* *	10	* *	36	* *		
0102	151	57.6 (109)	162	35.2 (111)	57	* *	45	* *	124	7.3 (75)		
0103	135	62.2 (113)	173	42.8 (118)	54	* *	38	* *	118	8.5 (88)		
0104	45	* *	51	* *	20	* *	13	* *	37	* *		
0250	54	* *	55	* *	26	* *	10	* *	42	* *		
0251	56	* *	54	* *	19	* *	14	* *	37	* *		
0252	43	* *	45	* *	14	* *	9	* *	40	* *		
0253	55	* *	67	* *	28	* *	14	* *	49	* *		
0401	196	62.2 (113)	215	35.3 (112)	83	* *	64	* *	155	4.5 (32)		
0402	138	51.4 (94)	161	33.5 (107)	58	* *	36	* *	135	4.4 (30)		
0403	45	* *	52	* *	16	* *	6	* *	36	* *		
0404	40	* *	54	* *	13	* *	11	* *	41	* *		
0501	186	53.8 (104)	211	27.5 (89)	65	* *	46	* *	144	12.5 (110)		
0502	0	* *	1	* *	0	* *	0	* *	1	* *		
0503	68	* *	76	* *	22	* *	17	* *	54	* *		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Smoking Indicators by Clinic

(1)

Page: 49
Run Date: 7 / 21 / 10

Clinic	Comparison of Smoking Indicators (2, 3)									
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal	Postpartum			
Clinic	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0504	205	53.2 (101)	222	32.9 (105)	80	* *	44	* *	157	5.7 (53)
0505	119	46.2 (75)	140	25.0 (75)	41	* *	39	* *	109	5.5 (50)
0506	67	* *	68	* *	20	* *	22	* *	53	* *
0650	380	48.9 (83)	444	29.1 (93)	170	30.6 (36)	237	19.8 (51)	440	7.3 (75)
0651	213	58.7 (111)	247	41.3 (116)	106	19.8 (52)	148	36.5 (83)	243	12.3 (109)
0652	89	* *	112	26.8 (86)	44	* *	67	* *	109	11.9 (108)
0653	142	52.8 (98)	164	32.9 (105)	69	* *	99	* *	162	20.4 (116)
0654	145	53.1 (100)	157	33.8 (108)	71	* *	93	* *	156	14.7 (114)
0655	123	61.8 (112)	141	44.7 (119)	70	* *	75	* *	139	10.1 (99)
0657	24	* *	27	* *	11	* *	17	* *	27	* *
0701	134	38.1 (53)	159	21.4 (58)	31	* *	27	* *	115	0.9 (1)
0702	26	* *	27	* *	10	* *	9	* *	22	* *
0703	81	* *	93	* *	24	* *	22	* *	68	* *
0704	20	* *	26	* *	11	* *	5	* *	20	* *
0705	52	* *	57	* *	23	* *	15	* *	37	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Smoking Indicators by Clinic

(1)

Page: 50
Run Date: 7 / 21 / 10

Clinic	Comparison of Smoking Indicators (2, 3)									
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal	Postpartum			
Clinic	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0850	381	44.4 (68)	443	20.3 (53)	147	38.1 (18)	359	20.1 (53)	432	5.3 (45)
0851	299	48.5 (82)	334	24.0 (67)	131	29.8 (41)	282	19.5 (47)	327	6.1 (59)
0901	625	53.3 (102)	673	31.2 (101)	261	33.0 (25)	243	22.2 (65)	574	5.9 (56)
0902	20	* *	27	* *	9	* *	6	* *	23	* *
1051	473	49.3 (86)	536	23.1 (65)	167	40.1 (17)	143	15.4 (31)	436	4.1 (25)
1101	593	39.8 (59)	744	21.5 (59)	190	31.1 (35)	489	23.1 (70)	741	6.1 (59)
1102	301	52.5 (97)	333	31.2 (101)	141	31.2 (34)	210	27.6 (79)	330	7.6 (80)
1106	37	* *	39	* *	18	* *	26	* *	39	* *
1250	372	39.0 (57)	418	26.6 (83)	130	23.1 (49)	284	15.8 (34)	416	4.8 (38)
1251	266	47.0 (80)	318	31.1 (100)	112	20.5 (51)	229	21.4 (60)	310	7.1 (70)
1252	231	34.6 (43)	230	19.6 (50)	65	* *	196	15.8 (34)	230	2.6 (9)
1253	253	49.4 (87)	346	29.8 (97)	109	24.8 (47)	218	21.6 (61)	344	5.8 (55)
1350	837	43.2 (66)	911	25.4 (77)	311	26.4 (42)	785	19.6 (49)	898	4.0 (24)
1351	62	* *	65	* *	22	* *	58	* *	65	* *
1352	150	54.0 (105)	149	35.6 (115)	76	* *	139	31.7 (82)	148	6.8 (68)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Smoking Indicators by Clinic

(1)

Page: 51
Run Date: 7 / 21 / 10

Clinic	Comparison of Smoking Indicators (2, 3)											
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household						
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal	Postpartum					
	Number	%	(Rank)	Number	%	(Rank)	Number	%	(Rank)	Number	%	(Rank)
1401	82	*	*	80	*	*	19	*	*	18	*	*
1402	58	*	*	72	*	*	15	*	*	17	*	*
1403	9	*	*	7	*	*	2	*	*	2	*	*
1404	10	*	*	5	*	*	0	*	*	3	*	*
1501	365	29.0	(29)	404	16.1	(40)	95	*	*	280	14.3	(25)
1503	432	35.6	(46)	494	22.1	(61)	137	31.4	(30)	330	15.5	(32)
1504	845	30.3	(33)	956	13.1	(28)	237	46.0	(6)	664	5.7	(2)
1505	294	29.6	(30)	338	13.9	(34)	80	*	*	234	7.3	(4)
1506	220	44.1	(67)	246	20.7	(55)	89	*	*	173	20.8	(56)
1507	211	45.0	(73)	249	22.1	(61)	92	*	*	169	16.6	(39)
1508	45	*	*	38	*	*	1	*	*	31	*	*
1509	242	29.8	(31)	265	19.2	(48)	66	*	*	172	8.7	(8)
1510	45	*	*	40	*	*	20	*	*	32	*	*
1511	226	51.3	(92)	256	35.5	(114)	107	22.4	(50)	181	9.9	(11)
1512	80	*	*	98	*	*	37	*	*	66	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Smoking Indicators by Clinic

(1)

Page: 52
Run Date: 7 / 21 / 10

Clinic	Comparison of Smoking Indicators (2, 3)											
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household						
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal	Postpartum					
	Number	%	(Rank)	Number	%	(Rank)	Number	%	(Rank)	Number	%	(Rank)
1701	250	52.0	(95)	281	30.2	(98)	77	*	*	42	*	*
1712	20	*	*	18	*	*	9	*	*	4	*	*
1950	259	46.7	(77)	289	24.6	(72)	106	32.1	(27)	170	26.5	(75)
1952	39	*	*	50	*	*	19	*	*	21	*	*
1953	97	*	*	115	29.6	(95)	49	*	*	59	*	*
1954	240	44.6	(70)	284	26.4	(82)	100	30.0	(40)	170	10.0	(12)
1955	55	*	*	56	*	*	24	*	*	31	*	*
1957	162	46.9	(79)	195	26.2	(81)	65	*	*	101	20.8	(56)
1959	60	*	*	60	*	*	27	*	*	34	*	*
2150	240	49.2	(84)	277	23.1	(65)	77	*	*	50	*	*
2151	100	58.0	(110)	114	32.5	(104)	37	*	*	27	*	*
2251	153	56.9	(108)	178	35.4	(113)	57	*	*	34	*	*
2252	99	*	*	98	*	*	32	*	*	20	*	*
2551	1,240	36.0	(49)	534	17.0	(43)	147	36.1	(20)	488	19.5	(47)
2554	33	*	*	31	*	*	11	*	*	12	*	*
										26	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Smoking Indicators by Clinic

(1)

Page: 53
Run Date: 7 / 21 / 10

Clinic	Comparison of Smoking Indicators (2, 3)									
	Smoking			Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household					
	3 Months Prior to Pregnancy	Last 3 Months of Pregnancy	Prenatal		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
2555	1,247	45.1 (74)	888	20.2 (52)	198	43.4 (12)	593	19.9 (52)	791	7.3 (75)
3001	344	34.3 (42)	373	19.3 (49)	92	* *	109	28.4 (81)	308	7.5 (79)
3002	89	* *	95	* *	22	* *	28	* *	78	* *
3003	1	* *	4	* *	0	* *	1	* *	4	* *
3004	64	* *	58	* *	3	* *	23	* *	44	* *
3007	7	* *	8	* *	1	* *	2	* *	7	* *
3009	2	* *	3	* *	0	* *	0	* *	3	* *
3010	0	* *	0	* *	0	* *	0	* *	0	* *
3201	184	46.7 (77)	221	25.8 (79)	67	* *	73	* *	179	6.1 (59)
3301	1,473	35.7 (48)	1,632	17.1 (44)	457	41.4 (13)	1,424	14.5 (27)	1,608	5.2 (44)
3302	87	* *	105	20.0 (51)	23	* *	86	* *	102	3.9 (22)
3304	29	* *	36	* *	1	* *	29	* *	36	* *
3305	26	* *	40	* *	8	* *	26	* *	39	* *
3306	58	* *	81	* *	22	* *	57	* *	80	* *
3308	13	* *	17	* *	8	* *	13	* *	17	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Smoking Indicators by Clinic

(1)

Page: 54
Run Date: 7 / 21 / 10

Clinic	Comparison of Smoking Indicators (2, 3)											
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household						
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum				
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)		
3309	8	* *	12	* *	2	* *	8	* *	12	* *		
3550	27	* *	32	* *	10	* *	10	* *	23	* *		
3551	7	* *	9	* *	3	* *	1	* *	6	* *		
3552	8	* *	13	* *	4	* *	0	* *	10	* *		
3701	367	41.1 (64)	408	24.5 (71)	134	26.1 (43)	216	24.1 (71)	404	6.2 (63)		
3805	1,181	49.5 (88)	1,251	28.4 (91)	520	31.7 (28)	1,105	20.9 (58)	1,232	6.3 (64)		
3950	723	38.2 (54)	780	22.1 (61)	245	35.1 (22)	502	14.9 (30)	779	4.7 (36)		
3952	5	* *	4	* *	2	* *	2	* *	4	* *		
3953	3	* *	2	* *	1	* *	1	* *	2	* *		
3954	23	* *	24	* *	7	* *	15	* *	23	* *		
4001	417	34.8 (44)	422	22.0 (60)	121	30.6 (36)	264	20.1 (53)	418	11.7 (107)		
4002	313	39.0 (57)	367	25.1 (76)	115	31.3 (32)	225	26.7 (77)	360	15.3 (115)		
4103	20	* *	32	* *	6	* *	14	* *	32	* *		
4106	244	29.9 (32)	312	13.8 (32)	62	* *	171	12.3 (16)	307	4.6 (34)		
4107	1,330	33.8 (40)	1,433	16.3 (42)	402	43.5 (11)	967	13.0 (21)	1,418	4.2 (27)		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Smoking Indicators by Clinic

(1)

Page: 55
Run Date: 7 / 21 / 10

Clinic	Comparison of Smoking Indicators (2, 3)									
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal	Postpartum			
Clinic	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4109	405	40.5 (63)	474	24.3 (69)	151	30.5 (38)	279	22.6 (68)	470	4.7 (36)
4110	626	26.8 (27)	675	15.7 (39)	150	25.3 (46)	452	13.3 (22)	667	3.4 (16)
4112	245	2.4 (1)	239	2.1 (1)	5	* *	169	6.5 (3)	237	3.8 (19)
4113	256	13.7 (8)	246	6.9 (6)	31	* *	158	1.9 (1)	243	1.6 (4)
4120	670	30.6 (34)	807	17.2 (45)	183	32.8 (26)	520	12.3 (16)	802	1.5 (3)
4160	606	25.7 (25)	703	13.4 (31)	137	43.8 (9)	475	7.8 (6)	693	2.0 (6)
4351	206	44.7 (71)	232	29.7 (96)	72	* *	68	* *	188	9.6 (96)
4352	58	* *	67	* *	21	* *	19	* *	51	* *
4353	169	40.2 (62)	187	20.9 (56)	49	* *	53	* *	151	6.0 (57)
4354	291	51.2 (91)	318	34.9 (110)	112	25.9 (44)	96	* *	246	11.0 (101)
4355	125	53.6 (103)	137	28.5 (92)	45	* *	42	* *	100	5.0 (41)
4356	205	44.4 (68)	222	26.6 (83)	59	* *	63	* *	161	5.0 (41)
4357	3	* *	4	* *	0	* *	0	* *	2	* *
4358	77	* *	75	* *	2	* *	31	* *	66	* *
4359	185	37.8 (52)	215	24.2 (68)	45	* *	54	* *	163	3.1 (13)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Smoking Indicators by Clinic

(1)

Page: 56
Run Date: 7 / 21 / 10

Clinic	Comparison of Smoking Indicators (2, 3)											
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household						
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum				
Clinic	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4360	52	* *	59	* *	14	* *	15	* *	42	* *		
4361	99	* *	101	41.6 (117)	47	* *	28	* *	78	* *		
4362	126	54.8 (107)	140	34.3 (109)	48	* *	28	* *	108	11.1 (102)		
4363	302	54.3 (106)	338	29.3 (94)	112	31.3 (32)	103	22.3 (66)	268	7.1 (70)		
4401	435	49.2 (84)	523	27.0 (87)	155	30.3 (39)	139	23.0 (69)	416	7.9 (82)		
4501	321	44.9 (72)	377	24.7 (73)	111	41.4 (13)	109	14.7 (28)	307	3.9 (22)		
4601	508	35.6 (46)	614	24.3 (69)	166	25.9 (44)	302	10.9 (14)	554	5.4 (48)		
4701	388	52.8 (98)	495	25.5 (78)	186	44.6 (8)	228	17.1 (41)	486	5.6 (51)		
5050	506	37.0 (51)	191	12.0 (23)	36	* *	243	16.5 (37)	169	4.1 (25)		
5051	985	38.8 (56)	443	13.3 (30)	96	* *	447	14.8 (29)	391	5.6 (51)		
5052	811	30.9 (35)	290	10.0 (15)	49	* *	400	12.3 (16)	262	3.1 (13)		
5101	672	36.0 (49)	874	18.3 (47)	194	40.7 (16)	362	14.4 (26)	808	4.6 (34)		
5202	230	50.9 (90)	251	27.5 (89)	84	* *	58	* *	185	3.2 (15)		
5203	61	* *	65	* *	19	* *	11	* *	42	* *		
5204	65	* *	70	* *	15	* *	10	* *	46	* *		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Smoking Indicators by Clinic

(1)

Page: 57
Run Date: 7 / 21 / 10

Clinic	Comparison of Smoking Indicators (2, 3)									
	Smoking			Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household					
	3 Months Prior to Pregnancy	Last 3 Months of Pregnancy	Prenatal		Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
5601	374	51.3 (92)	465	26.0 (80)	137	31.4 (30)	134	18.7 (46)	378	4.5 (32)
5801	787	47.5 (81)	873	26.7 (85)	324	31.5 (29)	455	19.6 (49)	850	7.9 (82)
5803	1	* *	17	* *	0	* *	1	* *	16	* *
5804	0	* *	7	* *	0	* *	0	* *	7	* *
6101	15	* *	16	* *	4	* *	11	* *	15	* *
6102	49	* *	64	* *	21	* *	40	* *	61	* *
6103	1,466	43.0 (65)	1,586	24.9 (74)	566	33.4 (24)	946	16.3 (36)	1,558	8.0 (84)
6104	31	* *	38	* *	11	* *	24	* *	36	* *
6105	14	* *	18	* *	8	* *	13	* *	18	* *
6106	1	* *	3	* *	0	* *	1	* *	3	* *
6301	1,471	31.0 (37)	1,829	14.5 (36)	358	40.8 (15)	766	12.7 (20)	1,691	4.9 (40)
6312	744	19.9 (19)	986	9.0 (14)	117	54.7 (3)	409	10.0 (12)	918	5.7 (53)
6316	416	38.7 (55)	560	15.4 (38)	135	46.7 (5)	224	8.9 (9)	516	2.5 (7)
6320	74	* *	90	* *	8	* *	35	* *	73	* *
6325	48	* *	55	* *	16	* *	21	* *	45	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Smoking Indicators by Clinic

(1)

Page: 58
Run Date: 7 / 21 / 10

Clinic	Comparison of Smoking Indicators (2, 3)									
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal	Postpartum			
Clinic	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
6330	178	30.9 (35)	188	14.4 (35)	40	* *	86	* *	169	8.3 (86)
6335	207	19.3 (18)	239	12.6 (26)	33	* *	99	* *	211	3.8 (19)
6340	48	* *	55	* *	8	* *	16	* *	49	* *
6345	16	* *	20	* *	8	* *	7	* *	18	* *
6350	80	* *	102	20.6 (54)	25	* *	44	* *	92	* *
7304	71	* *	82	* *	12	* *	22	* *	71	* *
7306	883	31.5 (38)	1,013	16.1 (40)	208	45.2 (7)	357	16.5 (37)	809	7.2 (73)
7308	72	* *	78	* *	29	* *	30	* *	67	* *
7310	19	* *	25	* *	10	* *	6	* *	20	* *
7401	801	52.4 (96)	808	30.4 (99)	338	37.0 (19)	331	21.1 (59)	730	8.8 (92)
7402	55	* *	52	* *	29	* *	19	* *	47	* *
7403	28	* *	25	* *	12	* *	5	* *	21	* *
7404	26	* *	23	* *	7	* *	9	* *	17	* *
7601	245	46.5 (76)	270	27.0 (87)	93	* *	95	* *	235	6.8 (68)
7850	311	49.5 (88)	357	32.2 (103)	137	24.8 (47)	177	24.3 (72)	355	5.4 (48)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Smoking Indicators by Clinic

(1)

Page: 59
Run Date: 7 / 21 / 10

Clinic	Comparison of Smoking Indicators (2, 3)											
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household						
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum				
Clinic	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
7851	53	* *	51	* *	29	* *	36	* *	51	* *		
7852	32	* *	39	* *	14	* *	16	* *	39	* *		
8150	14	* *	18	* *	1	* *	8	* *	17	* *		
8151	975	33.9 (41)	1,115	18.2 (46)	278	35.3 (21)	603	17.9 (43)	1,078	8.3 (86)		
8152	181	39.8 (59)	187	20.9 (56)	68	* *	102	37.3 (84)	184	13.0 (111)		
8153	8	* *	9	* *	4	* *	4	* *	9	* *		
8154	80	* *	92	* *	5	* *	39	* *	91	* *		
8155	18	* *	19	* *	1	* *	8	* *	19	* *		
8156	8	* *	8	* *	3	* *	5	* *	7	* *		
8157	13	* *	10	* *	2	* *	11	* *	10	* *		
8201	779	32.1 (39)	646	15.2 (37)	116	34.5 (23)	479	12.5 (19)	601	2.8 (11)		
8202	920	13.7 (8)	932	7.3 (8)	68	* *	510	8.2 (7)	867	3.8 (19)		
8203	94	* *	59	* *	7	* *	73	* *	53	* *		
8250	250	27.6 (28)	217	13.8 (32)	29	* *	246	22.4 (67)	214	4.2 (27)		
8255	869	39.9 (61)	349	12.0 (23)	71	* *	510	18.0 (44)	315	4.8 (38)		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Smoking Indicators by Clinic

(1)

Page: 60
Run Date: 7 / 21 / 10

Clinic	Comparison of Smoking Indicators (2, 3)									
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal	Postpartum			
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8257	166	6.0 (3)	145	6.9 (6)	5	* *	110	11.8 (15)	130	8.5 (88)
8258	497	35.4 (45)	371	22.6 (64)	83	* *	250	14.0 (24)	329	3.0 (12)
8260	157	19.1 (16)	172	12.2 (25)	25	* *	84	* *	160	6.3 (64)
8901	269	23.8 (21)	348	12.9 (27)	47	* *	144	25.7 (73)	296	11.5 (104)
8902	169	10.7 (4)	242	5.8 (3)	11	* *	95	* *	208	11.5 (104)
8903	360	17.5 (13)	444	10.1 (17)	53	* *	199	21.6 (61)	389	10.5 (100)
8904	256	14.5 (11)	331	8.5 (13)	28	* *	136	26.5 (75)	280	9.6 (96)
8905	778	14.5 (11)	1,040	10.6 (21)	88	* *	444	9.2 (10)	896	7.1 (70)
9004	3	* *	3	* *	0	* *	3	* *	3	* *
9012	10	* *	14	* *	0	* *	10	* *	12	* *
9013	12	* *	30	* *	1	* *	12	* *	30	* *
9015	9	* *	10	* *	0	* *	9	* *	10	* *
9016	21	* *	31	* *	1	* *	21	* *	31	* *
9017	8	* *	9	* *	0	* *	8	* *	9	* *
9018	19	* *	33	* *	2	* *	19	* *	31	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Smoking Indicators by Clinic

(1)

Page: 61
Run Date: 7 / 21 / 10

Clinic	Comparison of Smoking Indicators (2, 3)											
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household						
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum				
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)		
9019	10	* *	13	* *	0	* *	10	* *	12	* *		
9021	14	* *	20	* *	2	* *	14	* *	20	* *		
9040	3	* *	6	* *	1	* *	3	* *	6	* *		
9041	160	18.1 (15)	170	6.5 (4)	23	* *	34	* *	146	8.9 (93)		
9045	403	11.9 (5)	462	10.2 (18)	41	* *	225	15.6 (33)	431	7.2 (73)		
9050	626	12.6 (6)	749	7.6 (9)	72	* *	380	17.1 (41)	708	8.1 (85)		
9051	819	12.7 (7)	872	6.7 (5)	79	* *	361	7.5 (5)	827	2.5 (7)		
9053	758	25.1 (23)	1,017	11.5 (22)	158	43.7 (10)	489	21.7 (63)	971	13.5 (112)		
9055	32	* *	51	* *	1	* *	2	* *	21	* *		
9058	228	3.9 (2)	245	3.3 (2)	6	* *	135	17.0 (40)	234	6.0 (57)		
9066	676	13.9 (10)	878	7.9 (11)	78	* *	433	20.6 (55)	841	10.0 (98)		
9068	336	26.2 (26)	544	13.1 (28)	76	* *	249	27.7 (80)	520	14.6 (113)		
9069	790	20.4 (20)	1,022	7.6 (9)	143	57.3 (1)	539	26.0 (74)	1,001	8.9 (93)		
9070	600	18.0 (14)	770	8.4 (12)	87	* *	347	13.8 (23)	747	8.6 (90)		
9072	477	19.1 (16)	690	10.0 (15)	73	* *	366	18.3 (45)	665	11.3 (103)		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Smoking Indicators by Clinic

(1)

Page: 62
Run Date: 7 / 21 / 10

Clinic	Comparison of Smoking Indicators (2, 3)							
	Smoking			Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household			
	3 Months Prior to Pregnancy	Last 3 Months of Pregnancy	Prenatal		Number	% (Rank)	Number	% (Rank)
9074	504	24.2 (22)	648	10.2 (18)	104	54.8 (2)	337	27.0 (78)
9080	688	25.3 (24)	850	10.4 (20)	149	54.4 (4)	404	22.0 (64)
Unknown Clinics	8	* *	17	* *	0	* *	8	* *
Michigan	54,868	35.8	59,246	19.3	14,749	35.2	32,394	17.0
Nation (Prior Year)	999,199	23.1	856,953	14.2	146,933	35.5	655,095	18.9
							578,852	19.9

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Infant Health Indicators by Clinic

Page: 63
Run Date: 7 / 21 / 10

Clinic	Comparison of Infant Health Indicators (2, 3)									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)
0101	50	* * * *	54	* *	43	* *	52	* *		
0102	140	7.9 (60) 10.0 (99)	154	9.7 (29)	122	0.8 (1)	144	59.7 (70)		
0103	149	4.7 (4) 14.8 (115)	164	3.7 (1)	143	2.1 (14)	152	60.5 (66)		
0104	50	* * * *	50	* *	45	* *	50	* *		
0250	52	* * * *	53	* *	45	* *	52	* *		
0251	47	* * * *	52	* *	44	* *	47	* *		
0252	42	* * * *	40	* *	35	* *	42	* *		
0253	53	* * * *	57	* *	46	* *	53	* *		
0401	201	11.4 (102) 10.4 (103)	209	9.1 (13)	181	4.4 (87)	201	42.3 (104)		
0402	145	3.4 (1) 13.1 (114)	153	9.8 (34)	131	0.8 (1)	145	68.3 (24)		
0403	48	* * * *	48	* *	45	* *	48	* *		
0404	48	* * * *	50	* *	44	* *	48	* *		
0501	172	7.0 (37) 10.5 (107)	196	8.7 (9)	155	3.9 (79)	183	68.3 (24)		
0502	1	* * * *	1	* *	1	* *	1	* *		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Infant Health Indicators by Clinic

Page: 64
Run Date: 7 / 21 / 10

Clinic	Comparison of Infant Health Indicators (2, 3)									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)								
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
0503	69	*	*	75	*	59	*	70	*	*
0504	187	7.0 (37)	10.7 (109)	206	8.3 (8)	169	2.4 (24)	198	77.3 (6)	
0505	116	6.0 (18)	10.3 (102)	130	5.4 (2)	109	3.7 (74)	124	74.2 (12)	
0506	58	*	*	58	*	52	*	58	*	*
0650	413	5.3 (9)	6.3 (35)	433	12.7 (91)	360	2.2 (18)	414	65.7 (33)	
0651	229	6.1 (23)	5.7 (27)	239	11.3 (69)	203	4.4 (87)	229	55.9 (83)	
0652	98	*	*	109	9.2 (17)	89	*	100	61.0 (62)	
0653	151	7.9 (60)	7.9 (65)	154	11.0 (62)	134	3.7 (74)	151	64.9 (36)	
0654	143	7.7 (55)	9.8 (97)	148	7.4 (5)	130	3.1 (55)	143	72.0 (14)	
0655	125	8.0 (63)	5.6 (24)	136	9.6 (24)	112	2.7 (39)	125	74.4 (11)	
0657	26	*	*	25	*	25	*	26	*	*
0701	144	4.9 (8)	10.4 (103)	158	10.1 (43)	130	3.1 (55)	146	75.3 (8)	
0702	29	*	*	30	*	27	*	27	*	*
0703	88	*	*	88	*	78	*	88	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Infant Health Indicators by Clinic

Page: 65
Run Date: 7 / 21 / 10

Clinic	Comparison of Infant Health Indicators (2, 3)									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)								
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
0704	23	*	*	25	*	21	*	23	*	*
0705	53	*	*	55	*	47	*	53	*	*
0850	427	6.1 (23)	8.4 (79)	437	9.6 (24)	382	2.9 (46)	427	70.5 (17)	
0851	315	6.3 (28)	7.9 (65)	327	11.3 (69)	280	2.1 (14)	318	62.3 (55)	
0901	632	7.8 (56)	9.5 (92)	659	10.9 (59)	560	3.2 (60)	629	54.2 (85)	
0902	25	*	*	27	*	23	*	24	*	*
1051	480	5.4 (10)	9.6 (94)	514	12.3 (85)	419	1.4 (5)	483	83.6 (5)	
1101	693	7.6 (53)	5.8 (30)	727	9.8 (34)	622	3.5 (70)	692	46.2 (100)	
1102	313	5.8 (15)	8.3 (77)	320	9.7 (29)	280	1.8 (10)	312	50.0 (91)	
1106	38	*	*	38	*	32	*	38	*	*
1250	385	6.0 (18)	8.6 (83)	394	9.1 (13)	345	2.0 (13)	386	60.6 (64)	
1251	288	5.6 (12)	5.9 (32)	304	8.9 (11)	261	1.5 (6)	294	63.6 (44)	
1252	212	4.2 (2)	7.5 (56)	222	9.5 (21)	191	2.1 (14)	214	61.7 (57)	
1253	321	8.7 (81)	7.2 (47)	336	12.8 (92)	276	2.5 (28)	325	59.7 (70)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Infant Health Indicators by Clinic

Page: 66
Run Date: 7 / 21 / 10

Clinic	Comparison of Infant Health Indicators (2, 3)									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)
1350	870	7.5 (49) 8.0 (69)	896	9.4 (20)	786	2.8 (41)	875	49.6 (92)		
1351	64	* * * *	66		59	* *	64		*	*
1352	146	7.5 (49) 9.6 (94)	146	9.6 (24)	131	2.3 (22)	146	50.7 (90)		
1401	71	* * * *	78		64	* *	71		*	*
1402	68	* * * *	68		64	* *	68		*	*
1403	4	* * * *	6		4	* *	4		*	*
1404	6	* * * *	6		5	* *	6		*	*
1501	380	6.1 (23) 9.2 (89)	414	9.2 (17)	340	2.6 (32)	393	66.2 (31)		
1503	454	6.2 (27) 8.8 (86)	497	12.1 (82)	395	3.0 (52)	474	65.6 (34)		
1504	833	6.8 (36) 6.5 (40)	900	11.0 (62)	717	3.1 (55)	889	68.2 (26)		
1505	314	6.4 (29) 10.8 (110)	326	10.4 (48)	282	1.8 (10)	314	75.2 (9)		
1506	233	6.0 (18) 9.0 (88)	244	12.3 (85)	205	2.4 (24)	234	67.9 (27)		
1507	228	6.1 (23) 7.9 (65)	246	11.4 (74)	204	2.9 (46)	240	69.2 (20)		
1508	34	* * * *	53		31	* *	42		*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Infant Health Indicators by Clinic

Page: 67
Run Date: 7 / 21 / 10

Clinic	Comparison of Infant Health Indicators (2, 3)							
	Birthweight			Preterm (6)	Full Term LBW (7)		Ever Breastfed	
	Low (4)	High (5)	Number (Rank)		Number (Rank)	% (Rank)	Number (Rank)	% (Rank)
1509	252	7.1 (42) 7.5 (56)	269	11.2 (66)	220	3.2 (60)	257	61.1 (60)
1510	41	* * * *	42	*	36	* *	42	* *
1511	241	7.9 (60) 8.3 (77)	250	11.6 (76)	213	2.3 (22)	245	67.8 (28)
1512	89	* * * *	96	*	80	* *	90	* *
1701	257	7.0 (37) 10.9 (111)	279	7.2 (4)	234	3.8 (76)	255	63.1 (51)
1712	18	* * * *	18	*	17	* *	18	* *
1950	271	7.4 (48) 6.3 (35)	279	10.0 (41)	242	2.9 (46)	270	63.0 (52)
1952	49	* * * *	51	*	44	* *	49	* *
1953	106	10.4 (98) 5.7 (27)	106	12.3 (85)	92	* *	106	63.2 (46)
1954	239	8.8 (84) 7.5 (56)	272	11.8 (79)	208	4.3 (85)	260	63.8 (42)
1955	45	* * * *	56	*	42	* *	52	* *
1957	181	8.3 (73) 11.0 (112)	190	14.2 (102)	153	3.3 (64)	182	67.6 (29)
1959	58	* * * *	59	*	52	* *	58	* *
2150	262	6.5 (30) 9.5 (92)	272	10.3 (46)	235	3.8 (76)	262	61.1 (60)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Infant Health Indicators by Clinic

Page: 68
Run Date: 7 / 21 / 10

Clinic	Comparison of Infant Health Indicators (2, 3)									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)								
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
2151	105	9.5 (88)	10.5 (107)	113	9.7 (29)	95	* *	104	47.1 (99)	
2251	161	7.5 (49)	8.1 (73)	163	9.8 (34)	144	2.8 (41)	161	59.0 (72)	
2252	88	*	*	94	*	74	*	88	*	
2551	499	9.6 (90)	3.8 (8)	521	13.8 (101)	426	2.6 (32)	502	59.8 (69)	
2554	30	*	*	31	*	29	*	30	*	
2555	830	7.8 (56)	7.5 (56)	883	11.2 (66)	723	3.0 (52)	839	69.1 (21)	
3001	330	13.6 (110)	2.7 (2)	355	13.0 (95)	288	9.0 (113)	331	47.4 (96)	
3002	83	*	*	91	*	70	*	83	*	
3003	4	*	*	4	*	4	*	4	*	
3004	50	*	*	53	*	42	*	50	*	
3007	7	*	*	9	*	7	*	6	*	
3009	5	*	*	6	*	5	*	5	*	
3010	0	*	*	1	*	0	*	0	*	
3201	191	4.7 (4)	7.9 (65)	209	11.0 (62)	168	2.4 (24)	193	53.9 (86)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Infant Health Indicators by Clinic

Page: 69
Run Date: 7 / 21 / 10

Clinic	Comparison of Infant Health Indicators (2, 3)									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)								
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
3301	1,562	6.7 (32)	7.4 (54)	1,622	10.2 (45)	1,398	2.5 (28)	1,580	62.7 (54)	
3302	96	*	*	104	14.4 (105)	83	*	97	*	
3304	35	*	*	35	*	30	*	35	*	
3305	37	*	*	38	*	36	*	38	*	
3306	79	*	*	83	*	70	*	80	*	
3308	17	*	*	18	*	16	*	17	*	
3309	11	*	*	11	*	9	*	11	*	
3550	29	*	*	29	*	27	*	29	*	
3551	9	*	*	9	*	9	*	9	*	
3552	14	*	*	14	*	14	*	14	*	
3701	389	5.7 (13)	6.7 (42)	401	12.2 (83)	341	2.1 (14)	389	63.2 (46)	
3805	1,207	7.6 (53)	7.2 (47)	1,221	9.7 (29)	1,071	3.4 (68)	1,210	59.0 (72)	
3950	749	8.4 (75)	8.0 (69)	767	10.4 (48)	663	2.7 (39)	750	58.7 (74)	
3952	4	*	*	4	*	4	*	4	*	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Infant Health Indicators by Clinic

Page: 70
Run Date: 7 / 21 / 10

Clinic	Comparison of Infant Health Indicators (2, 3)									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)								
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
3953	2	*	*	2	*	2	*	2	*	
3954	22	*	*	22	*	19	*	22	*	
4001	371	13.7 (111)	3.5 (6)	402	17.4 (113)	299	4.7 (92)	386	56.2 (82)	
4002	342	8.2 (69)	5.6 (24)	354	14.7 (106)	292	3.1 (55)	345	63.2 (46)	
4103	30	*	*	29	*	25	*	30	*	
4106	296	9.5 (88)	5.1 (21)	306	10.8 (54)	262	4.6 (91)	297	60.3 (67)	
4107	1,349	8.2 (69)	7.3 (51)	1,391	11.1 (65)	1,197	3.2 (60)	1,357	64.3 (39)	
4109	448	8.5 (77)	10.0 (99)	460	13.0 (95)	387	2.8 (41)	445	60.2 (68)	
4110	630	6.7 (32)	7.6 (60)	651	10.0 (41)	564	3.2 (60)	638	62.1 (56)	
4112	231	4.8 (7)	10.4 (103)	234	9.8 (34)	206	1.5 (6)	233	63.5 (45)	
4113	235	7.2 (44)	6.8 (43)	241	10.8 (54)	210	1.9 (12)	234	65.4 (35)	
4120	757	6.7 (32)	8.2 (75)	784	9.8 (34)	679	1.6 (8)	765	63.8 (42)	
4160	658	8.2 (69)	8.1 (73)	679	9.6 (24)	589	4.8 (93)	663	61.7 (57)	
4351	218	7.3 (46)	11.0 (112)	226	6.2 (3)	201	4.5 (90)	219	70.3 (18)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Infant Health Indicators by Clinic

Page: 71
Run Date: 7 / 21 / 10

Clinic	Comparison of Infant Health Indicators (2, 3)									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)								
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
4352	66	*	*	68	*	62	*	66	*	*
4353	169	6.5 (30)	6.5 (40)	179	9.5 (21)	149	3.4 (68)	171	63.2 (46)	
4354	301	4.7 (4)	8.0 (69)	303	10.6 (51)	263	0.8 (1)	302	62.9 (53)	
4355	121	9.9 (94)	9.9 (98)	134	14.2 (102)	103	5.8 (103)	121	68.6 (22)	
4356	206	7.8 (56)	8.7 (84)	213	12.2 (83)	180	3.9 (79)	205	61.5 (59)	
4357	3	*	*	3	*	3	*	3	*	*
4358	71	*	*	74	*	60	*	73	*	*
4359	192	4.2 (2)	7.3 (51)	210	9.0 (12)	176	1.7 (9)	193	72.0 (14)	
4360	54	*	*	56	*	47	*	55	*	*
4361	90	*	*	97	*	82	*	90	*	*
4362	127	7.1 (42)	9.4 (91)	134	9.7 (29)	115	4.3 (85)	128	77.3 (6)	
4363	309	8.1 (66)	10.4 (103)	324	9.6 (24)	277	2.2 (18)	310	71.6 (16)	
4401	482	8.9 (85)	6.4 (39)	505	10.3 (46)	428	3.5 (70)	480	65.8 (32)	
4501	349	6.0 (18)	7.4 (54)	357	9.2 (17)	312	2.9 (46)	349	57.0 (81)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Infant Health Indicators by Clinic

Page: 72
Run Date: 7 / 21 / 10

Clinic	Comparison of Infant Health Indicators (2, 3)									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)								
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
4601	520	8.1 (66)	6.9 (45)	622	9.5 (21)	465	3.9 (79)	590	63.2 (46)	
4701	469	7.0 (37)	10.0 (99)	482	9.1 (13)	426	2.6 (32)	468	64.5 (38)	
5050	183	8.7 (81)	3.3 (3)	188	12.8 (92)	160	5.0 (95)	183	96.7 (1)	
5051	424	8.3 (73)	8.0 (69)	443	11.3 (69)	372	3.5 (70)	424	89.4 (3)	
5052	281	8.5 (77)	4.6 (18)	293	10.6 (51)	252	2.8 (41)	282	89.7 (2)	
5101	830	7.5 (49)	5.5 (23)	869	10.8 (54)	735	2.6 (32)	842	64.0 (41)	
5202	242	5.8 (15)	8.7 (84)	245	13.5 (99)	208	2.4 (24)	242	68.6 (22)	
5203	60	*	*	64	*	53	*	61	*	
5204	71	*	*	71	*	66	*	71	*	
5601	439	8.2 (69)	8.9 (87)	457	10.9 (59)	389	3.3 (64)	437	72.1 (13)	
5801	827	8.0 (63)	9.3 (90)	855	12.9 (94)	719	2.6 (32)	827	57.1 (80)	
5803	18	*	*	19	*	18	*	18	*	
5804	6	*	*	6	*	4	*	6	*	
6101	15	*	*	16	*	15	*	16	*	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Infant Health Indicators by Clinic

Page: 73
Run Date: 7 / 21 / 10

Clinic	Comparison of Infant Health Indicators (2, 3)									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)								
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
6102	61	*	*	61	*	52	*	61	*	*
6103	1,480	8.4 (75)	7.2 (47)	1,529	12.5 (89)	1,281	3.3 (64)	1,484	48.2 (95)	
6104	34	*	*	37	*	28	*	34	*	*
6105	18	*	*	18	*	16	*	18	*	*
6106	3	*	*	3	*	3	*	3	*	*
6301	1,750	8.7 (81)	7.3 (51)	1,804	11.3 (69)	1,540	3.8 (76)	1,761	47.2 (97)	
6312	955	9.9 (94)	6.0 (33)	983	12.4 (88)	836	3.0 (52)	958	57.8 (76)	
6316	535	6.0 (18)	9.7 (96)	549	8.2 (7)	490	2.2 (18)	535	57.2 (79)	
6320	80	*	*	89	*	69	*	85	*	*
6325	52	*	*	53	*	43	*	52	*	*
6330	176	11.4 (102)	6.3 (35)	183	18.0 (115)	140	2.9 (46)	180	39.4 (105)	
6335	223	7.2 (44)	8.5 (80)	234	9.8 (34)	199	4.0 (82)	227	42.7 (103)	
6340	54	*	*	53	*	48	*	55	*	*
6345	19	*	*	19	*	16	*	19	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Infant Health Indicators by Clinic

Page: 74
Run Date: 7 / 21 / 10

Clinic	Comparison of Infant Health Indicators (2, 3)									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)								
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
6350	92	*	*	102	7.8 (6)	85	*	94	*	*
7304	73	*	*	78	*	64	*	74	*	*
7306	937	8.9 (85)	6.8 (43)	980	10.4 (48)	831	4.1 (83)	936	47.2 (97)	
7308	75	*	*	77	*	63	*	75	*	*
7310	25	*	*	25	*	25	*	25	*	*
7401	771	9.3 (87)	7.1 (46)	792	12.6 (90)	668	2.5 (28)	772	52.7 (87)	
7402	52	*	*	52	*	46	*	52	*	*
7403	25	*	*	25	*	21	*	25	*	*
7404	22	*	*	22	*	20	*	22	*	*
7601	260	5.4 (10)	4.2 (12)	264	11.7 (78)	226	2.2 (18)	257	57.6 (78)	
7850	341	7.3 (46)	7.6 (60)	353	9.9 (40)	307	2.6 (32)	343	60.6 (64)	
7851	50	*	*	51	*	44	*	50	*	*
7852	36	*	*	37	*	30	*	36	*	*
8150	15	*	*	18	*	12	*	15	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Infant Health Indicators by Clinic

Page: 75
Run Date: 7 / 21 / 10

Clinic	Comparison of Infant Health Indicators (2, 3)									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)								
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
8151	1,052	8.1 (66)	8.2 (75)	1,083	11.8 (79)	921	3.5 (70)	1,056	66.3 (30)	
8152	181	9.9 (94)	7.7 (62)	186	10.8 (54)	160	4.4 (87)	182	54.4 (84)	
8153	7	*	*	7	*	6	*	7	*	
8154	88	*	*	91	*	81	*	88	*	
8155	17	*	*	18	*	17	*	17	*	
8156	7	*	*	8	*	7	*	8	*	
8157	9	*	*	10	*	9	*	9	*	
8201	598	8.0 (63)	8.5 (80)	643	10.7 (53)	531	2.8 (41)	602	64.8 (37)	
8202	781	7.0 (37)	6.3 (35)	914	11.2 (66)	687	2.9 (46)	871	69.3 (19)	
8203	53	*	*	59	*	49	*	52	*	
8250	214	9.8 (93)	5.6 (24)	216	8.8 (10)	191	5.8 (103)	215	57.7 (77)	
8255	309	7.8 (56)	6.1 (34)	337	10.1 (43)	274	3.3 (64)	311	88.4 (4)	
8257	120	6.7 (32)	4.2 (12)	137	10.9 (59)	106	0.9 (4)	130	74.6 (10)	
8258	328	5.8 (15)	8.5 (80)	364	9.1 (13)	294	3.1 (55)	330	58.2 (75)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Infant Health Indicators by Clinic

Page: 76
Run Date: 7 / 21 / 10

Clinic	Comparison of Infant Health Indicators (2, 3)									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)								
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
8260	163	8.6 (80)	3.7 (7)	166	10.8 (54)	145	6.2 (109)	165	64.2 (40)	
8901	311	13.2 (108)	3.9 (10)	333	15.3 (110)	258	5.4 (99)	312	25.3 (116)	
8902	230	11.7 (105)	4.8 (19)	237	18.1 (116)	188	5.3 (97)	230	30.4 (112)	
8903	409	13.2 (108)	3.4 (4)	427	15.5 (112)	338	5.6 (101)	411	28.0 (115)	
8904	309	13.9 (112)	4.5 (14)	324	14.8 (108)	261	5.7 (102)	311	44.4 (102)	
8905	957	9.6 (90)	7.2 (47)	1,002	11.6 (76)	827	5.4 (99)	967	32.8 (109)	
9004	3	*	*	3	*	2	*	3	*	
9012	12	*	*	13	*	9	*	13	*	
9013	29	*	*	29	*	23	*	29	*	
9015	9	*	*	10	*	6	*	9	*	
9016	30	*	*	30	*	22	*	30	*	
9017	9	*	*	9	*	9	*	9	*	
9018	32	*	*	33	*	26	*	32	*	
9019	13	*	*	13	*	12	*	13	*	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Infant Health Indicators by Clinic

Page: 77
Run Date: 7 / 21 / 10

Clinic	Comparison of Infant Health Indicators (2, 3)									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)	Number (%) (Rank)							
9021	18 * * * *	20 * *	16 * *	18 * *						
9040	5 * * * *	6 * *	5 * *	5 * *						
9041	155 10.3 (97) 4.5 (14)	159 11.3 (69)	134 5.2 (96)	153 32.7 (110)						
9045	434 9.7 (92) 7.8 (64)	448 15.2 (109)	361 4.2 (84)	437 46.2 (100)						
9050	703 10.5 (99) 5.7 (27)	722 15.4 (111)	586 5.3 (97)	707 30.3 (113)						
9051	778 5.7 (13) 7.7 (62)	842 11.9 (81)	671 2.5 (28)	829 52.1 (88)						
9053	954 11.6 (104) 4.5 (14)	1,002 14.7 (106)	806 4.8 (93)	954 37.2 (106)						
9055	49 * * * *	51 * *	46 * *	49 * *						
9058	223 8.5 (77) 5.8 (30)	226 11.5 (75)	195 2.6 (32)	225 60.9 (63)						
9066	832 13.1 (107) 3.4 (4)	866 14.3 (104)	705 7.1 (112)	837 28.1 (114)						
9068	515 14.6 (113) 3.9 (10)	524 17.4 (113)	412 6.1 (107)	516 35.9 (107)						
9069	966 11.3 (101) 4.5 (14)	1,004 13.1 (97)	830 5.8 (103)	989 48.4 (94)						
9070	717 10.7 (100) 3.8 (8)	755 13.1 (97)	613 5.9 (106)	732 34.2 (108)						
9072	632 16.1 (115) 4.9 (20)	676 18.9 (118)	506 6.7 (111)	637 30.8 (111)						

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Infant Health Indicators by Clinic

Page: 78
Run Date: 7 / 21 / 10

Clinic	Comparison of Infant Health Indicators (2, 3)									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)
9074	622	15.1 (114)	2.6 (1)		634	18.8 (117)	497	6.4 (110)	623	49.3 (93)
9080	786	11.7 (105)	5.3 (22)		821	13.6 (100)	676	6.1 (107)	795	51.3 (89)
Unknown Clinics	10	*	*	*	28	*	10	*	11	*
Michigan	55,121	8.4	7.1		57,774	11.5	48,299	3.5	55,786	57.2
Nation (Prior Year)	1,085,134	8.2	6.6		1,025,278	11.5	890,981	3.6	952,980	65.4

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

(1)

Page: 79
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Racial and Ethnic Distribution ⁽²⁾							
		White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %	
0101	165	97.0	0.6	0.6	0.0	0.0	1.2	0.6	
0102	505	95.8	0.6	1.4	0.4	0.2	1.6	0.0	
0103	512	98.4	0.2	0.6	0.0	0.2	0.6	0.0	
0104	151	94.0	0.7	1.3	0.0	0.0	4.0	0.0	
0250	186	89.8	0.5	1.1	1.1	0.0	7.5	0.0	
0251	165	93.3	0.0	0.0	3.6	1.2	1.8	0.0	
0252	155	68.4	1.3	1.3	1.9	2.6	24.5	0.0	
0253	208	81.7	0.0	0.5	12.0	0.5	5.3	0.0	
0401	653	95.9	0.5	0.9	0.0	0.5	2.0	0.3	
0402	540	89.1	0.4	0.4	0.2	0.6	9.4	0.0	
0403	150	98.7	0.0	0.0	0.7	0.0	0.7	0.0	
0404	169	95.9	0.6	0.6	0.0	0.0	3.0	0.0	
0501	647	96.9	0.0	1.1	0.5	0.2	1.4	0.0	
0502	4	*	*	*	*	*	*	*	
0503	196	92.9	0.0	2.0	0.5	0.0	4.1	0.5	
0504	679	91.8	0.6	1.0	2.5	1.0	3.1	0.0	
0505	453	86.5	1.3	2.0	0.7	0.0	9.5	0.0	
0506	240	97.1	0.0	1.3	0.4	0.4	0.8	0.0	
0650	1,319	91.1	2.6	2.0	0.8	0.9	2.3	0.2	
0651	779	95.6	0.3	1.3	0.1	0.5	1.5	0.6	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

(1)

Page: 80
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Racial and Ethnic Distribution ⁽²⁾							
		White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %	
0652	401	96.8	0.0	0.5	0.0	0.0	1.7	1.0	
0653	516	99.0	0.0	0.6	0.0	0.0	0.4	0.0	
0654	461	96.1	0.7	1.1	0.7	0.0	1.3	0.2	
0655	500	97.4	0.2	0.4	0.0	0.4	1.6	0.0	
0657	83	*	*	*	*	*	*	*	
0701	511	92.2	1.0	0.4	0.8	4.9	0.6	0.2	
0702	111	94.6	0.0	0.9	0.0	0.0	3.6	0.9	
0703	304	96.4	0.0	1.3	0.7	0.7	1.0	0.0	
0704	88	*	*	*	*	*	*	*	
0705	215	98.1	0.0	0.9	0.0	0.5	0.5	0.0	
0850	1,400	92.0	3.3	3.0	0.0	0.6	0.9	0.2	
0851	1,078	95.8	0.4	2.9	0.0	0.4	0.3	0.3	
0901	2,163	84.6	2.8	8.6	0.0	0.5	3.4	0.1	
0902	86	*	*	*	*	*	*	*	
1051	1,774	93.2	0.5	3.7	0.5	0.2	1.5	0.4	
1055	2	*	*	*	*	*	*	*	
1101	2,303	38.0	54.2	3.9	0.2	0.9	2.0	0.9	
1102	1,083	80.7	11.3	4.3	0.2	0.7	2.5	0.3	
1106	158	89.9	0.6	3.8	0.6	0.6	3.8	0.6	
1250	1,269	86.1	0.3	9.9	0.0	2.3	1.2	0.2	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

(1)

Page: 81
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Racial and Ethnic Distribution ⁽²⁾							
		White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %	
1251	1,059	95.8	0.5	1.8	0.0	0.1	0.8	1.0	
1252	928	65.1	0.6	32.7	0.1	0.2	1.2	0.1	
1253	1,042	88.9	3.9	4.6	0.0	0.1	2.1	0.4	
1350	3,064	64.8	20.5	8.5	0.3	1.7	3.6	0.6	
1351	182	90.1	2.2	4.4	0.0	0.0	2.2	1.1	
1352	534	66.3	24.7	4.1	0.0	0.7	3.7	0.4	
1401	289	93.8	0.0	4.2	0.0	0.0	1.7	0.3	
1402	198	57.6	0.0	21.2	11.1	0.5	9.6	0.0	
1403	32	*	*	*	*	*	*	*	
1404	28	*	*	*	*	*	*	*	
1501	1,430	47.6	6.7	40.3	0.3	0.1	2.4	2.7	
1502	14	*	*	*	*	*	*	*	
1503	1,699	58.1	5.6	31.2	0.4	0.6	1.9	2.2	
1504	3,148	46.3	4.2	40.8	0.1	5.8	1.9	0.8	
1505	1,023	87.0	1.9	7.3	0.0	1.2	2.1	0.6	
1506	756	80.3	1.1	15.5	0.3	0.1	2.1	0.7	
1507	811	93.5	0.1	3.6	0.0	0.2	2.2	0.4	
1508	265	0.8	0.0	94.7	0.0	0.0	0.0	4.5	
1509	858	51.6	2.0	42.3	0.0	0.3	1.9	1.9	
1510	240	84.6	4.6	8.8	0.0	0.0	2.1	0.0	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

(1)

Page: 82
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Racial and Ethnic Distribution ⁽²⁾							
		White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %	
1511	743	94.3	1.2	2.8	0.0	0.0	1.1	0.5	
1512	345	96.5	0.6	1.2	0.3	0.0	1.2	0.3	
1701	777	63.1	0.5	0.9	23.8	1.5	9.8	0.4	
1712	97	*	*	*	*	*	*	*	
1950	836	95.3	0.1	3.7	0.0	0.0	0.6	0.2	
1952	192	96.4	0.0	2.6	0.0	0.0	1.0	0.0	
1953	410	95.9	0.2	2.7	0.0	0.5	0.7	0.0	
1954	785	91.7	0.0	7.1	0.0	0.1	0.8	0.3	
1955	174	92.5	0.6	5.2	0.0	0.0	1.7	0.0	
1957	550	92.0	0.7	4.7	0.5	0.4	1.1	0.5	
1959	239	89.5	4.2	3.8	0.0	0.4	2.1	0.0	
1963	36	*	*	*	*	*	*	*	
2150	838	94.7	0.0	0.7	0.5	0.1	3.5	0.5	
2151	362	95.9	0.0	1.1	0.3	0.3	1.4	1.1	
2251	576	95.0	0.5	0.9	0.2	0.3	2.6	0.5	
2252	288	94.1	0.3	1.7	0.0	0.0	3.8	0.0	
2551	4,588	30.4	67.6	1.2	0.0	0.2	0.6	0.0	
2552	8	*	*	*	*	*	*	*	
2554	138	92.0	5.8	0.7	0.0	0.7	0.7	0.0	
2555	5,168	76.4	19.8	2.6	0.1	0.4	0.7	0.1	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

(1)

Page: 83
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Racial and Ethnic Distribution ⁽²⁾							
		White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %	
3001	1,211	12.7	68.4	16.9	0.0	0.7	1.3	0.0	
3002	279	19.4	55.6	21.9	0.4	0.0	2.9	0.0	
3003	11	*	*	*	*	*	*	*	
3004	229	3.9	0.0	94.8	0.0	0.0	0.9	0.4	
3007	44	*	*	*	*	*	*	*	
3009	19	*	*	*	*	*	*	*	
3010	4	*	*	*	*	*	*	*	
3201	696	95.5	0.1	2.9	0.0	0.6	0.6	0.3	
3301	5,149	49.7	27.5	15.1	0.1	3.7	3.3	0.6	
3302	366	24.9	49.5	14.8	0.0	6.6	4.1	0.3	
3303	17	*	*	*	*	*	*	*	
3304	132	39.4	15.9	6.1	0.0	36.4	2.3	0.0	
3305	130	73.8	9.2	10.8	0.0	4.6	1.5	0.0	
3306	214	93.0	0.0	2.8	0.0	0.9	1.4	1.9	
3308	65	*	*	*	*	*	*	*	
3309	43	*	*	*	*	*	*	*	
3311	13	*	*	*	*	*	*	*	
3550	94	*	*	*	*	*	*	*	
3551	38	*	*	*	*	*	*	*	
3552	54	*	*	*	*	*	*	*	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

(1)

Page: 84
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Racial and Ethnic Distribution (2)							
		White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %	
3701	1,127	90.8	0.5	6.7	0.2	0.4	1.3	0.1	
3702	133	94.0	0.8	4.5	0.0	0.8	0.0	0.0	
3703	57	*	*	*	*	*	*	*	
3704	56	*	*	*	*	*	*	*	
3805	4,106	77.4	12.6	5.9	0.1	0.8	3.1	0.1	
3950	2,597	67.3	18.0	7.0	0.1	1.7	5.7	0.2	
3952	52	*	*	*	*	*	*	*	
3953	40	*	*	*	*	*	*	*	
3954	139	81.3	4.3	3.6	0.7	5.8	4.3	0.0	
4001	1,339	29.1	55.9	10.8	0.2	0.9	2.9	0.1	
4002	1,143	53.6	27.0	13.6	0.2	1.2	4.1	0.3	
4103	127	11.8	75.6	6.3	0.0	2.4	3.9	0.0	
4106	1,236	10.3	51.1	34.7	0.2	0.5	2.6	0.6	
4107	4,166	54.2	27.1	12.2	0.3	2.0	4.0	0.1	
4109	1,460	91.8	0.5	5.0	0.2	0.5	1.8	0.2	
4110	2,126	39.6	9.7	46.3	0.2	0.5	3.4	0.2	
4111	1	*	*	*	*	*	*	*	
4112	823	0.9	0.5	98.1	0.0	0.0	0.4	0.2	
4113	1,001	8.1	17.2	70.6	0.0	1.5	2.6	0.0	
4120	2,511	61.1	8.8	24.6	0.2	2.4	2.8	0.1	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

(1)

Page: 85
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Racial and Ethnic Distribution ⁽²⁾							
		White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %	
4160	2,340	45.3	33.2	10.6	0.2	6.6	4.1	0.1	
4351	794	94.2	0.3	3.0	0.0	0.1	1.9	0.5	
4352	226	76.1	15.5	3.1	0.4	0.4	4.4	0.0	
4353	660	75.5	0.2	21.8	0.0	0.0	2.0	0.6	
4354	959	92.2	2.9	1.1	0.1	0.7	2.9	0.0	
4355	479	81.4	0.0	6.3	1.3	0.2	9.8	1.0	
4356	692	86.3	0.0	8.5	0.7	0.9	3.3	0.3	
4357	23	*	*	*	*	*	*	*	
4358	291	3.1	0.0	95.9	0.0	0.0	0.0	1.0	
4359	658	72.5	0.2	25.1	0.3	0.3	1.1	0.6	
4360	219	93.6	0.0	6.4	0.0	0.0	0.0	0.0	
4361	310	92.6	2.3	0.6	0.0	0.3	3.9	0.3	
4362	420	94.5	0.0	1.2	0.7	0.0	3.3	0.2	
4363	1,152	94.7	0.1	2.3	0.1	0.4	2.1	0.3	
4401	1,619	93.0	0.5	3.0	0.1	0.3	3.0	0.1	
4501	1,190	95.7	0.3	2.7	0.0	0.1	1.1	0.1	
4601	1,972	77.5	1.4	18.1	0.0	0.3	1.7	1.0	
4701	1,562	93.3	0.5	3.8	0.1	1.3	0.6	0.4	
5050	1,542	71.5	22.8	1.7	0.3	1.2	2.5	0.1	
5051	3,217	67.6	22.8	4.3	0.1	1.9	3.1	0.2	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

(1)

Page: 86
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Racial and Ethnic Distribution ⁽²⁾							
		White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %	
5052	2,279	67.1	21.0	2.0	0.1	7.6	2.0	0.2	
5054	2	*	*	*	*	*	*	*	
5101	1,987	70.7	8.9	15.2	0.1	0.9	3.8	0.5	
5202	712	91.7	0.1	1.0	1.8	0.3	5.1	0.0	
5203	240	90.0	2.9	0.8	0.4	0.8	5.0	0.0	
5204	223	89.7	2.2	0.4	0.4	0.4	6.7	0.0	
5205	40	*	*	*	*	*	*	*	
5601	1,433	96.0	1.1	0.8	0.0	0.5	1.3	0.2	
5801	2,726	88.6	3.7	5.2	0.2	0.4	1.3	0.6	
5803	26	*	*	*	*	*	*	*	
5804	11	*	*	*	*	*	*	*	
6101	47	*	*	*	*	*	*	*	
6102	170	94.7	1.2	1.8	0.0	0.0	2.4	0.0	
6103	5,217	61.9	26.3	7.7	0.2	0.6	3.2	0.2	
6104	81	*	*	*	*	*	*	*	
6105	45	*	*	*	*	*	*	*	
6106	5	*	*	*	*	*	*	*	
6301	5,784	48.2	27.7	19.3	0.1	2.7	1.8	0.2	
6312	3,212	40.4	54.4	1.3	0.2	1.8	1.7	0.2	
6316	1,639	75.2	12.7	8.1	0.2	2.4	1.0	0.3	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

(1)

Page: 87
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Racial and Ethnic Distribution ⁽²⁾							
		White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %	
6320	315	70.8	24.4	1.0	0.0	3.2	0.6	0.0	
6325	188	95.2	1.1	1.6	0.0	0.0	2.1	0.0	
6330	542	31.0	49.1	17.3	0.2	0.7	1.7	0.0	
6335	727	79.6	13.8	2.8	0.0	2.9	1.0	0.0	
6340	224	45.5	14.7	35.3	0.0	3.6	0.9	0.0	
6345	61	*	*	*	*	*	*	*	
6350	335	63.9	31.0	2.1	0.0	1.2	1.8	0.0	
7304	284	95.8	0.4	3.5	0.0	0.4	0.0	0.0	
7306	3,284	50.9	34.3	12.0	0.1	0.5	2.0	0.2	
7308	237	94.5	0.4	3.4	0.0	0.0	1.7	0.0	
7310	34	*	*	*	*	*	*	*	
7401	2,494	84.2	5.8	5.1	0.1	0.6	4.1	0.1	
7402	293	94.2	0.0	1.7	0.0	0.0	4.1	0.0	
7403	114	85.1	0.0	13.2	1.8	0.0	0.0	0.0	
7404	112	92.0	0.9	2.7	0.0	0.0	4.5	0.0	
7601	896	94.3	0.1	4.0	0.0	0.1	0.9	0.6	
7850	1,174	94.0	0.1	3.3	0.0	0.1	2.0	0.5	
7851	219	93.6	0.5	4.1	0.0	0.0	1.8	0.0	
7852	133	97.7	0.8	0.0	0.0	0.0	1.5	0.0	
8150	83	*	*	*	*	*	*	*	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

(1)

Page: 88
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Racial and Ethnic Distribution ⁽²⁾							
		White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %	
8151	3,400	43.5	41.3	9.5	0.4	2.2	3.0	0.1	
8152	602	43.9	42.2	3.2	0.2	1.0	9.6	0.0	
8153	41	*	*	*	*	*	*	*	
8154	377	35.0	17.0	11.4	0.0	35.5	1.1	0.0	
8155	107	56.1	28.0	8.4	0.0	6.5	0.9	0.0	
8156	51	*	*	*	*	*	*	*	
8157	27	*	*	*	*	*	*	*	
8201	2,654	62.7	24.9	4.9	0.3	2.6	4.4	0.0	
8202	3,575	90.2	5.3	2.4	0.1	1.0	0.9	0.1	
8203	324	72.8	21.6	2.8	0.0	0.3	2.2	0.3	
8250	355	45.1	46.8	2.8	0.0	2.8	2.5	0.0	
8255	3,185	70.1	22.6	4.3	0.0	1.2	1.6	0.1	
8257	596	51.2	22.8	0.2	0.0	24.8	1.0	0.0	
8258	1,811	64.4	27.5	4.5	0.1	0.7	2.7	0.2	
8260	490	47.3	31.8	1.8	0.0	16.7	2.2	0.0	
8901	1,162	1.1	96.2	0.6	0.0	0.7	1.4	0.0	
8902	716	3.9	93.3	0.8	0.0	0.1	1.8	0.0	
8903	1,401	2.4	95.3	0.5	0.1	0.7	0.9	0.0	
8904	1,045	5.3	79.7	0.7	0.1	12.3	1.9	0.0	
8905	2,879	30.1	31.2	27.7	0.2	6.3	4.3	0.2	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

(1)

Page: 89
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Racial and Ethnic Distribution ⁽²⁾							
		White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %	
9004	336	0.9	98.5	0.3	0.3	0.0	0.0	0.0	
9010	36	*	*	*	*	*	*	*	
9012	16	*	*	*	*	*	*	*	
9013	34	*	*	*	*	*	*	*	
9015	11	*	*	*	*	*	*	*	
9016	39	*	*	*	*	*	*	*	
9017	11	*	*	*	*	*	*	*	
9018	34	*	*	*	*	*	*	*	
9019	14	*	*	*	*	*	*	*	
9021	22	*	*	*	*	*	*	*	
9023	80	*	*	*	*	*	*	*	
9040	273	0.7	98.2	0.0	0.0	0.7	0.4	0.0	
9041	1,013	1.1	98.0	0.4	0.0	0.1	0.4	0.0	
9045	1,510	9.3	49.6	39.5	0.0	0.3	1.1	0.1	
9050	2,354	13.2	55.5	29.1	0.1	1.4	0.6	0.1	
9051	3,133	8.1	10.2	81.0	0.0	0.1	0.6	0.0	
9053	3,449	1.6	95.8	0.6	0.1	0.8	1.1	0.1	
9054	83	*	*	*	*	*	*	*	
9055	113	0.9	95.6	1.8	0.0	0.9	0.9	0.0	
9058	632	75.9	18.0	0.6	0.0	4.9	0.5	0.0	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 3A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Racial and Ethnic Distribution by Clinic
Combined 3 Years

(1)

Page: 90
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Racial and Ethnic Distribution ⁽²⁾							
		White, Not Hispanic %	Black, Not Hispanic %	Hispanic %	American Indian/Alaska Native %	Asian/Pacific Islander %	Multiple Races %	All Other/Unknown %	
9066	2,919	1.0	97.6	0.1	0.1	0.1	1.1	0.0	
9068	1,725	1.9	96.2	0.5	0.0	0.6	0.5	0.2	
9069	2,745	1.5	96.8	0.4	0.1	0.3	0.9	0.0	
9070	2,504	1.3	91.6	0.0	0.1	6.1	0.9	0.0	
9072	2,026	1.7	94.8	0.2	0.1	2.2	0.8	0.1	
9074	1,907	2.2	94.0	0.6	0.1	0.2	2.8	0.1	
9079	548	0.5	98.2	0.2	0.0	0.5	0.5	0.0	
9080	2,644	24.2	71.4	2.2	0.1	0.2	1.7	0.2	
Unknown Clinics	463	5.8	2.6	22.9	2.4	2.4	62.2	1.7	
Michigan	203,605	56.9	27.7	10.8	0.3	1.6	2.4	0.3	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Combined 3 Years

Page: 91
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Age Distribution (2)							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
0101	165	0.0	6.1	12.1	61.8	18.2	1.2	0.6	
0102	505	0.2	5.0	11.7	68.1	13.7	1.4	0.0	
0103	512	0.2	4.7	14.6	63.5	15.0	2.0	0.0	
0104	151	0.0	6.6	12.6	66.9	13.9	0.0	0.0	
0250	186	0.0	10.8	13.4	61.8	13.4	0.5	0.0	
0251	165	0.0	4.2	9.7	70.9	12.7	2.4	0.0	
0252	155	0.6	1.9	12.9	61.9	21.9	0.6	0.0	
0253	208	0.0	7.7	8.7	63.0	19.2	1.4	0.0	
0401	653	0.2	5.8	14.5	65.4	13.0	1.1	0.0	
0402	540	0.0	4.3	14.1	61.1	18.7	1.9	0.0	
0403	150	0.0	2.0	8.0	72.7	16.0	1.3	0.0	
0404	169	0.0	3.6	8.3	65.1	21.9	1.2	0.0	
0501	647	0.0	3.9	14.8	66.5	14.4	0.5	0.0	
0502	4	*	*	*	*	*	*	*	
0503	196	0.0	1.5	16.3	64.3	15.8	1.5	0.5	
0504	679	0.0	4.3	13.0	64.8	16.9	1.0	0.0	
0505	453	0.0	3.5	10.8	67.5	16.8	1.3	0.0	
0506	240	0.0	9.2	18.8	59.6	10.8	1.7	0.0	
0650	1,319	0.0	6.7	12.7	65.7	13.5	0.9	0.4	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Age Distribution by Clinic
Combined 3 Years

(1)

Page: 92
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Age Distribution ⁽²⁾							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
0651	779	0.3	6.7	17.3	62.8	12.3	0.6	0.0	
0652	401	0.2	3.5	10.7	66.8	16.5	2.0	0.2	
0653	516	0.4	8.3	13.8	64.1	12.2	1.0	0.2	
0654	461	0.4	7.6	12.4	63.6	14.8	1.1	0.2	
0655	500	0.0	7.8	13.2	64.4	14.2	0.4	0.0	
0657	83	*	*	*	*	*	*	*	*
0701	511	0.0	2.3	10.2	66.5	18.0	2.9	0.0	
0702	111	0.0	6.3	9.0	70.3	13.5	0.9	0.0	
0703	304	0.0	2.3	10.9	64.1	22.0	0.7	0.0	
0704	88	*	*	*	*	*	*	*	*
0705	215	0.0	3.3	7.4	61.4	23.7	4.2	0.0	
0850	1,400	0.2	5.4	13.2	64.4	16.1	0.7	0.0	
0851	1,078	0.2	7.5	14.4	64.4	12.3	1.2	0.0	
0901	2,163	0.3	6.7	13.7	61.3	17.0	1.0	0.0	
0902	86	*	*	*	*	*	*	*	*
1051	1,774	0.3	5.1	11.4	62.5	19.4	1.2	0.0	
1055	2	*	*	*	*	*	*	*	*
1101	2,303	0.3	10.2	14.4	58.5	15.1	1.4	0.1	
1102	1,083	0.8	6.6	14.5	61.2	15.6	1.0	0.2	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Combined 3 Years

Page: 93
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Age Distribution (2)							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
1106	158	0.0	5.7	12.7	56.3	22.8	2.5	0.0	
1250	1,269	0.5	6.7	13.9	66.2	12.0	0.6	0.1	
1251	1,059	0.3	5.2	13.2	66.4	14.1	0.8	0.0	
1252	928	0.1	6.3	14.3	60.2	17.9	1.2	0.0	
1253	1,042	0.6	6.3	15.8	62.2	14.3	0.8	0.0	
1350	3,064	0.5	7.9	15.7	60.5	14.8	0.5	0.1	
1351	182	0.0	7.7	13.2	61.0	18.1	0.0	0.0	
1352	534	0.7	9.2	13.3	61.4	13.9	1.5	0.0	
1401	289	0.0	5.2	9.3	65.4	17.6	1.7	0.7	
1402	198	0.0	6.6	12.1	59.1	20.2	2.0	0.0	
1403	32	*	*	*	*	*	*	*	*
1404	28	*	*	*	*	*	*	*	*
1501	1,430	0.3	8.5	15.7	57.8	16.3	1.4	0.1	
1502	14	*	*	*	*	*	*	*	*
1503	1,699	0.2	6.6	12.7	61.9	17.3	1.2	0.1	
1504	3,148	0.4	7.3	12.3	60.3	18.6	1.0	0.0	
1505	1,023	0.0	4.5	12.1	61.9	20.0	1.5	0.0	
1506	756	0.1	7.3	14.0	60.4	17.1	1.1	0.0	
1507	811	0.1	5.3	11.7	65.7	16.4	0.6	0.1	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Combined 3 Years

Page: 94
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Age Distribution (2)							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
1508	265	0.0	10.9	7.9	58.5	20.0	2.6	0.0	
1509	858	0.2	7.7	14.7	56.9	19.0	1.5	0.0	
1510	240	0.0	7.5	14.6	66.3	10.4	1.3	0.0	
1511	743	0.1	5.7	14.7	63.4	14.9	1.2	0.0	
1512	345	0.0	4.3	12.5	60.6	21.2	1.4	0.0	
1701	777	0.1	4.9	11.1	66.2	17.4	0.3	0.1	
1712	97	*	*	*	*	*	*	*	*
1950	836	0.2	7.7	17.8	61.2	11.7	0.8	0.5	
1952	192	0.0	2.1	14.1	63.0	19.8	1.0	0.0	
1953	410	0.2	5.6	13.9	64.9	14.6	0.5	0.2	
1954	785	0.3	4.5	13.5	67.5	13.1	1.0	0.1	
1955	174	0.0	5.2	15.5	65.5	12.6	1.1	0.0	
1957	550	0.2	4.9	13.1	62.5	17.8	1.5	0.0	
1959	239	0.0	4.6	15.5	61.9	16.7	1.3	0.0	
1963	36	*	*	*	*	*	*	*	*
2150	838	0.1	3.7	10.9	67.4	17.1	0.8	0.0	
2151	362	0.6	5.8	11.6	66.0	15.2	0.6	0.3	
2251	576	0.3	4.3	12.0	64.4	17.7	1.2	0.0	
2252	288	1.0	6.9	12.8	63.5	14.9	0.7	0.0	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Combined 3 Years

Page: 95
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Age Distribution (2)							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
2551	4,588	0.7	10.6	16.8	57.5	13.1	1.2	0.0	
2552	8	*	*	*	*	*	*	*	*
2554	138	0.7	11.6	15.9	58.7	11.6	1.4	0.0	
2555	5,168	0.3	5.4	13.0	63.5	16.8	0.9	0.1	
3001	1,211	1.5	14.0	15.6	53.2	13.5	2.1	0.1	
3002	279	0.7	10.0	13.6	62.4	12.9	0.4	0.0	
3003	11	*	*	*	*	*	*	*	*
3004	229	0.0	5.2	13.1	55.0	24.9	1.7	0.0	
3007	44	*	*	*	*	*	*	*	*
3009	19	*	*	*	*	*	*	*	*
3010	4	*	*	*	*	*	*	*	*
3201	696	0.0	3.3	9.8	63.6	22.1	1.1	0.0	
3301	5,149	0.4	6.8	13.7	61.3	16.7	1.1	0.0	
3302	366	0.5	7.4	10.7	61.5	18.9	1.1	0.0	
3303	17	*	*	*	*	*	*	*	*
3304	132	0.0	0.0	4.5	46.2	47.7	0.8	0.8	
3305	130	0.0	4.6	7.7	63.1	21.5	2.3	0.8	
3306	214	0.0	1.9	10.7	65.9	21.0	0.0	0.5	
3308	65	*	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Combined 3 Years

Page: 96
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Age Distribution (2)							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
3309	43	*	*	*	*	*	*	*	*
3311	13	*	*	*	*	*	*	*	*
3550	94	*	*	*	*	*	*	*	*
3551	38	*	*	*	*	*	*	*	*
3552	54	*	*	*	*	*	*	*	*
3701	1,127	0.1	5.5	14.2	67.6	11.6	1.0	0.0	
3702	133	1.5	4.5	13.5	65.4	14.3	0.8	0.0	
3703	57	*	*	*	*	*	*	*	*
3704	56	*	*	*	*	*	*	*	*
3805	4,106	0.5	7.7	15.2	61.1	14.0	0.9	0.7	
3950	2,597	0.5	5.8	12.6	63.1	17.0	0.8	0.1	
3952	52	*	*	*	*	*	*	*	*
3953	40	*	*	*	*	*	*	*	*
3954	139	0.7	1.4	10.1	61.9	25.2	0.7	0.0	
4001	1,339	0.5	10.6	15.8	58.0	13.6	1.5	0.0	
4002	1,143	0.8	7.0	14.6	62.3	14.4	0.9	0.0	
4103	127	0.0	9.4	11.0	52.0	25.2	2.4	0.0	
4106	1,236	0.5	8.7	13.8	59.5	15.9	1.2	0.3	
4107	4,166	0.6	7.7	13.7	59.6	16.5	1.3	0.6	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Combined 3 Years

Page: 97
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Age Distribution (2)							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
4109	1,460	0.1	5.7	12.3	62.8	17.5	1.5	0.1	
4110	2,126	0.5	8.1	13.0	59.2	18.7	0.5	0.0	
4111	1	*	*	*	*	*	*	*	*
4112	823	0.2	6.1	8.0	57.8	26.0	1.6	0.2	
4113	1,001	0.5	7.3	13.6	56.5	21.0	0.8	0.3	
4120	2,511	0.1	5.9	11.5	63.4	18.0	1.0	0.1	
4160	2,340	0.4	5.1	11.7	61.9	19.8	1.0	0.1	
4351	794	0.1	7.9	15.5	61.7	14.4	0.4	0.0	
4352	226	0.4	6.6	14.6	62.8	14.6	0.9	0.0	
4353	660	0.3	7.7	14.8	61.1	14.8	1.2	0.0	
4354	959	0.0	5.1	14.9	64.9	13.8	1.3	0.1	
4355	479	0.2	6.5	15.0	62.8	14.2	1.3	0.0	
4356	692	0.3	6.1	14.2	64.9	14.0	0.6	0.0	
4357	23	*	*	*	*	*	*	*	
4358	291	1.0	7.2	10.3	56.0	22.7	2.7	0.0	
4359	658	0.0	6.1	13.8	62.0	17.2	0.9	0.0	
4360	219	0.0	7.8	11.0	60.3	20.1	0.9	0.0	
4361	310	3.2	8.4	15.2	60.3	11.6	1.3	0.0	
4362	420	0.0	5.2	13.3	66.7	14.5	0.2	0.0	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Combined 3 Years

Page: 98
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Age Distribution (2)							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
4363	1,152	0.1	5.6	12.3	68.2	12.8	1.0	0.0	
4401	1,619	0.0	3.6	13.6	64.2	17.4	1.2	0.0	
4501	1,190	0.3	4.7	12.8	62.4	18.8	1.0	0.1	
4601	1,972	0.2	6.3	16.0	62.2	14.5	0.8	0.0	
4701	1,562	0.1	4.4	11.6	63.4	19.0	1.5	0.0	
5050	1,542	0.1	4.5	10.4	59.6	24.1	1.3	0.0	
5051	3,217	0.2	3.6	10.7	63.0	21.0	1.4	0.1	
5052	2,279	0.3	4.7	10.0	59.3	24.0	1.7	0.0	
5054	2	*	*	*	*	*	*	*	*
5101	1,987	0.2	3.5	10.7	60.8	22.8	2.0	0.1	
5202	712	0.0	2.8	11.0	67.4	17.8	0.8	0.1	
5203	240	0.0	3.3	10.8	65.0	20.0	0.8	0.0	
5204	223	0.0	2.7	12.1	70.4	13.9	0.9	0.0	
5205	40	*	*	*	*	*	*	*	*
5601	1,433	0.1	5.0	12.8	66.7	14.7	0.7	0.0	
5801	2,726	0.2	7.0	14.3	61.7	15.8	0.9	0.0	
5803	26	*	*	*	*	*	*	*	*
5804	11	*	*	*	*	*	*	*	*
6101	47	*	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Combined 3 Years

Page: 99
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Age Distribution (2)							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
6102	170	0.0	7.6	10.0	63.5	18.8	0.0	0.0	
6103	5,217	0.3	7.3	14.8	63.5	13.2	0.8	0.0	
6104	81	*	*	*	*	*	*	*	*
6105	45	*	*	*	*	*	*	*	*
6106	5	*	*	*	*	*	*	*	*
6301	5,784	0.3	6.2	12.7	60.7	18.8	1.3	0.0	
6312	3,212	0.3	3.7	9.6	56.6	27.5	2.3	0.1	
6316	1,639	0.1	4.1	7.6	63.2	23.0	2.0	0.0	
6320	315	0.0	2.2	7.9	58.1	31.4	0.3	0.0	
6325	188	0.0	4.3	12.8	63.8	18.6	0.5	0.0	
6330	542	0.4	8.3	18.3	54.1	17.9	1.1	0.0	
6335	727	0.3	3.0	8.5	60.4	24.5	3.3	0.0	
6340	224	0.0	3.6	10.3	66.1	18.8	1.3	0.0	
6345	61	*	*	*	*	*	*	*	*
6350	335	0.0	9.9	13.1	57.9	17.6	1.2	0.3	
7304	284	0.0	6.3	10.9	63.7	18.3	0.7	0.0	
7306	3,284	0.4	7.5	14.0	60.1	16.8	1.1	0.1	
7308	237	0.0	3.4	9.7	66.7	19.0	1.3	0.0	
7310	34	*	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Combined 3 Years

Page: 100
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Age Distribution (2)							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
7401	2,494	0.5	6.2	14.6	64.0	13.8	0.9	0.0	
7402	293	0.3	1.7	12.3	61.4	23.2	1.0	0.0	
7403	114	0.0	3.5	14.9	60.5	20.2	0.9	0.0	
7404	112	0.0	8.9	11.6	64.3	14.3	0.9	0.0	
7601	896	0.2	4.1	12.2	67.2	15.5	0.7	0.1	
7850	1,174	0.2	7.2	15.2	62.4	14.1	1.1	0.0	
7851	219	0.0	5.9	15.1	56.6	20.5	1.8	0.0	
7852	133	0.8	1.5	14.3	67.7	15.0	0.8	0.0	
8150	83	*	*	*	*	*	*	*	*
8151	3,400	0.0	2.8	7.8	64.9	23.2	1.1	0.1	
8152	602	0.7	18.1	40.9	40.2	0.0	0.0	0.2	
8153	41	*	*	*	*	*	*	*	*
8154	377	0.0	1.6	2.4	58.1	37.1	0.8	0.0	
8155	107	0.0	3.7	4.7	54.2	35.5	1.9	0.0	
8156	51	*	*	*	*	*	*	*	*
8157	27	*	*	*	*	*	*	*	*
8201	2,654	0.1	4.6	13.6	59.6	20.7	1.3	0.1	
8202	3,575	0.1	3.0	7.2	60.1	26.5	2.9	0.2	
8203	324	0.3	4.3	13.3	62.0	18.2	1.9	0.0	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Combined 3 Years

Page: 101
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Age Distribution (2)							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
8250	355	0.0	4.2	15.2	58.9	20.3	1.4	0.0	
8255	3,185	0.3	5.4	12.8	62.3	16.6	0.9	1.7	
8257	596	0.3	4.0	9.6	62.8	20.5	2.9	0.0	
8258	1,811	0.2	6.2	11.9	59.6	20.9	1.2	0.0	
8260	490	0.2	4.1	10.0	58.2	24.5	2.7	0.4	
8901	1,162	1.1	11.2	14.5	55.8	16.3	1.1	0.0	
8902	716	0.8	4.9	15.8	56.0	21.4	1.1	0.0	
8903	1,401	1.0	11.5	14.7	52.5	19.1	1.3	0.0	
8904	1,045	0.5	7.0	12.2	56.7	22.4	1.1	0.1	
8905	2,879	0.6	6.5	11.4	55.2	24.2	2.0	0.1	
9004	336	0.9	13.4	13.4	51.5	19.0	1.8	0.0	
9010	36	*	*	*	*	*	*	*	*
9012	16	*	*	*	*	*	*	*	*
9013	34	*	*	*	*	*	*	*	*
9015	11	*	*	*	*	*	*	*	*
9016	39	*	*	*	*	*	*	*	*
9017	11	*	*	*	*	*	*	*	*
9018	34	*	*	*	*	*	*	*	*
9019	14	*	*	*	*	*	*	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Combined 3 Years

Page: 102
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Age Distribution (2)							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
9021	22	*	*	*	*	*	*	*	*
9023	80	*	*	*	*	*	*	*	*
9040	273	1.5	9.5	16.8	46.2	25.3	0.7	0.0	
9041	1,013	0.7	12.6	16.2	47.8	21.3	1.2	0.2	
9045	1,510	2.3	19.1	13.3	48.5	15.4	1.3	0.1	
9050	2,354	0.5	7.1	12.7	53.1	24.1	2.4	0.1	
9051	3,133	0.3	7.8	10.7	55.9	23.5	1.8	0.1	
9053	3,449	0.4	7.5	15.3	55.4	19.6	1.7	0.1	
9054	83	*	*	*	*	*	*	*	*
9055	113	1.8	6.2	20.4	51.3	19.5	0.9	0.0	
9058	632	0.2	2.4	5.5	56.5	31.5	4.0	0.0	
9066	2,919	0.3	8.5	15.8	53.8	19.2	1.5	0.9	
9068	1,725	0.5	9.0	13.8	56.7	18.5	1.5	0.1	
9069	2,745	0.5	7.8	15.2	54.4	20.6	1.4	0.0	
9070	2,504	0.5	9.0	14.6	54.6	19.6	1.6	0.0	
9072	2,026	0.6	10.0	17.7	53.1	17.3	1.2	0.0	
9074	1,907	0.3	7.4	16.3	57.5	17.0	1.5	0.0	
9079	548	1.1	6.8	9.7	50.7	30.3	1.5	0.0	
9080	2,644	0.3	6.0	12.7	56.5	22.7	1.8	0.0	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 4A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Age Distribution by Clinic
Combined 3 Years

Page: 103
Run Date: 7 / 21 / 10

Clinic	Number Included in Analysis	Comparison of Age Distribution (2)							
		< 15 Years %	15 - 17 Years %	18 - 19 Years %	20 - 29 Years %	30 - 39 Years %	>= 40 Years %	Unknown %	
Unknown Clinics	463	0.0	0.6	7.8	71.7	19.0	0.6	0.2	
Michigan	203,605	0.3	6.6	13.2	60.4	18.1	1.3	0.1	

(1) Reporting period is January 1 through December 31.

(2) Excludes data with errors.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Page: 104
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Health Indicators (2, 3)											
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Gestational Diabetes During Pregnancy	Hypertension During Pregnancy	
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester	Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0101	152	3.3 (40)	54.6 (108)	132	22.7 (69)	53.0 (135)	14	* *	24	* *	39	* * 39
0102	473	5.5 (146)	53.9 (95)	402	28.1 (118)	47.5 (72)	55	* *	84	* *	123	7.3 (80) 124
0103	493	5.1 (122)	48.7 (20)	427	30.0 (132)	50.1 (97)	52	* *	94	* *	122	8.2 (95) 122
0104	140	5.7 (152)	51.4 (50)	121	20.7 (49)	50.4 (104)	12	* *	30	* *	38	* * 38
0250	174	2.3 (8)	56.3 (124)	142	65.5 (167)	18.3 (4)	11	* *	41	* *	42	* * 42
0251	158	3.2 (34)	51.9 (57)	139	50.4 (162)	25.9 (8)	12	* *	23	* *	36	* * 36
0252	147	4.1 (70)	52.4 (67)	124	27.4 (110)	48.4 (79)	11	* *	38	* *	40	* * 40
0253	195	5.1 (122)	52.8 (73)	164	58.5 (164)	22.6 (6)	11	* *	33	* *	51	* * 51
0401	622	4.8 (106)	52.6 (71)	552	18.7 (22)	55.3 (157)	77	* *	156	23.1 (18)	151	4.6 (26) 151
0402	526	3.0 (27)	54.6 (108)	473	21.1 (55)	50.5 (107)	45	* *	137	21.9 (16)	132	4.5 (23) 132
0403	146	6.2 (162)	52.1 (60)	123	19.5 (32)	55.3 (157)	10	* *	33	* *	35	* * 35
0404	163	5.5 (146)	51.5 (53)	144	21.5 (61)	50.7 (111)	23	* *	42	* *	42	* * 42
0501	500	3.6 (45)	50.6 (42)	435	25.5 (98)	45.1 (51)	50	* *	107	60.7 (99)	147	6.1 (60) 147
0502	3	* * * *	* *	2	* * * *	* *	0	* *	1	* *	1	* * 1
0503	163	5.5 (146)	49.7 (25)	144	27.8 (114)	41.0 (26)	20	* *	32	* *	52	* * 52
0504	504	4.4 (87)	51.4 (50)	431	25.1 (92)	46.2 (60)	57	* *	91	* *	156	3.8 (9) 156
0505	349	5.4 (141)	55.9 (121)	295	43.7 (161)	34.9 (15)	50	* *	72	* *	105	8.6 (99) 106
												17.0(115)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Page: 105
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Health Indicators (2, 3)											
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Gestational Diabetes During Pregnancy	Hypertension During Pregnancy	
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester	Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0506	221	4.5 (96)	56.1 (123)	181	37.0 (150)	44.2 (45)	19	* *	43	* *	51	* *
0650	1,259	4.4 (87)	53.1 (81)	1,052	28.2 (121)	47.0 (68)	144	19.4 (5)	238	25.6 (29)	454	3.1 (3)
0651	733	5.6 (151)	54.0 (96)	617	29.3 (128)	44.7 (49)	71	* *	173	32.9 (61)	260	3.5 (8)
0652	369	2.2 (6)	58.3 (146)	323	31.3 (135)	46.1 (56)	40	* *	91	* *	123	9.8 (105)
0653	488	4.7 (103)	50.4 (37)	413	36.1 (149)	44.6 (46)	41	* *	87	* *	162	5.6 (48)
0654	436	3.9 (54)	55.0 (114)	352	21.6 (63)	49.4 (90)	43	* *	100	28.0 (40)	150	4.0 (10)
0655	471	5.3 (133)	50.3 (34)	424	21.2 (56)	52.1 (126)	47	* *	179	12.8 (2)	146	15.1 (116)
0657	81	* *	* *	68	* *	* *	14	* *	20	* *	28	* *
0701	452	4.4 (87)	43.1 (5)	385	24.7 (84)	43.9 (42)	49	* *	109	29.4 (47)	115	4.3 (17)
0702	102	2.9 (24)	52.0 (58)	90	* *	* *	13	* *	8	* *	23	* *
0703	286	5.9 (157)	52.4 (67)	253	19.0 (26)	56.5 (164)	31	* *	54	* *	66	* *
0704	83	* *	* *	75	* *	* *	11	* *	20	* *	20	* *
0705	184	1.6 (3)	50.5 (38)	163	24.5 (82)	42.3 (32)	20	* *	40	* *	40	* *
0850	1,323	3.5 (44)	53.1 (81)	1,114	18.2 (13)	55.0 (155)	175	18.9 (4)	331	24.2 (23)	640	5.0 (35)
0851	1,025	5.2 (130)	50.3 (34)	886	28.7 (125)	48.0 (77)	123	23.6 (10)	197	29.4 (47)	484	4.1 (13)
0901	2,018	5.4 (141)	54.2 (103)	1,787	24.8 (86)	50.2 (99)	229	28.8 (16)	547	24.9 (26)	564	8.7 (100)
0902	80	* *	* *	76	* *	* *	11	* *	32	* *	23	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

(1)

Page: 106
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Health Indicators (2, 3)											
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Gestational Diabetes During Pregnancy	Hypertension During Pregnancy	
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester	Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1051	1,661	4.2 (77)	50.0 (28)	1,427	20.0 (37)	51.8 (122)	153	30.7 (23)	434	20.3 (10)	425	4.9 (32)
1055	1	* * * *	* *	1	* * * *	* *	0	* *	0	* * *	0	* *
1101	2,045	4.4 (87)	56.3 (124)	1,718	26.9 (108)	48.4 (79)	362	50.3 (67)	693	48.5 (94)	849	5.8 (53)
1102	958	5.7 (152)	50.5 (38)	815	19.0 (26)	54.2 (147)	134	29.1 (18)	373	34.0 (66)	404	4.5 (23)
1106	142	7.7 (170)	47.9 (15)	118	16.9 (7)	58.5 (166)	27	* *	53	* *	47	* *
1250	1,208	4.9 (116)	52.5 (69)	1,030	28.1 (118)	46.1 (56)	150	20.7 (6)	152	26.3 (33)	491	6.5 (70)
1251	992	4.8 (106)	54.2 (103)	844	28.3 (122)	46.1 (56)	138	13.8 (3)	245	31.4 (54)	381	4.2 (15)
1252	837	3.3 (40)	56.4 (127)	572	31.1 (134)	41.1 (28)	112	35.7 (39)	139	35.3 (67)	309	12.9 (115)
1253	934	5.0 (118)	50.6 (42)	681	33.3 (142)	41.6 (29)	141	31.2 (24)	184	35.3 (67)	408	6.1 (60)
1350	2,877	4.2 (77)	55.2 (115)	2,489	20.0 (37)	53.0 (135)	487	35.5 (37)	480	44.2 (86)	1,322	5.6 (48)
1351	168	2.4 (12)	53.0 (78)	147	19.7 (35)	53.1 (137)	33	* *	33	* *	91	* *
1352	509	2.4 (12)	55.2 (115)	437	20.4 (43)	53.3 (140)	68	* *	60	* *	220	5.9 (57)
1401	265	2.6 (20)	62.3 (167)	220	27.7 (113)	49.1 (88)	18	* *	53	* *	60	* *
1402	177	2.3 (8)	60.5 (159)	159	16.4 (5)	50.9 (114)	23	* *	50	* *	58	* *
1403	32	* * * *	* *	27	* * * *	* *	0	* *	10	* *	4	* *
1404	23	* * * *	* *	8	* * * *	* *	3	* *	3	* *	6	* *
1501	1,170	3.6 (45)	56.5 (129)	916	40.2 (155)	37.3 (18)	152	39.5 (51)	199	42.2 (82)	479	8.1 (94)
											477	6.9 (18)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Page: 107
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Health Indicators (2, 3)											
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Gestational Diabetes During Pregnancy	Hypertension During Pregnancy	
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester	Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1502	13	* *	* *	11	* *	* *	1	* *	4	* *	0	* *
1503	1,325	5.1 (122)	56.4 (127)	973	61.7 (165)	22.7 (7)	193	33.7 (31)	226	36.7 (71)	572	5.4 (45)
1504	2,729	4.1 (70)	52.3 (64)	2,098	41.6 (159)	35.2 (16)	423	37.4 (44)	552	31.3 (53)	1,183	8.3 (97)
1505	949	4.0 (61)	47.9 (15)	842	21.3 (58)	46.7 (67)	132	37.1 (42)	243	30.9 (51)	420	3.3 (5)
1506	642	4.8 (106)	52.2 (62)	535	27.9 (116)	48.4 (79)	87	* *	94	* *	285	7.7 (87)
1507	729	4.0 (61)	51.4 (50)	599	19.7 (35)	55.8 (162)	90	* *	154	32.5 (59)	308	5.8 (53)
1508	177	6.8 (166)	53.7 (92)	114	66.7 (168)	13.2 (1)	26	* *	18	* *	48	* *
1509	723	3.2 (34)	56.7 (133)	485	40.2 (155)	34.8 (14)	78	* *	89	* *	312	9.6 (104)
1510	218	4.1 (70)	52.8 (73)	177	33.3 (142)	39.0 (20)	24	* *	39	* *	55	* *
1511	683	3.1 (31)	58.9 (150)	562	18.5 (19)	55.9 (163)	86	* *	116	31.0 (52)	311	7.1 (76)
1512	302	4.3 (83)	57.9 (143)	229	24.0 (76)	41.9 (31)	44	* *	69	* *	131	5.3 (40)
1701	749	5.3 (133)	51.5 (53)	641	20.1 (39)	54.8 (153)	60	* *	145	12.4 (1)	192	5.2 (38)
1712	92	* *	* *	81	* *	* *	5	* *	8	* *	15	* *
1950	805	5.3 (133)	52.8 (73)	692	23.8 (74)	50.4 (104)	83	* *	226	16.4 (6)	298	4.0 (10)
1952	189	5.3 (133)	54.0 (96)	171	18.1 (11)	48.5 (83)	25	* *	77	* *	60	* *
1953	399	5.3 (133)	54.1 (100)	346	23.7 (72)	50.0 (93)	43	* *	125	21.6 (15)	123	11.4 (112)
1954	694	6.3 (164)	53.3 (88)	563	26.3 (105)	50.3 (102)	81	* *	196	20.4 (12)	292	5.1 (37)
											294	9.2 (49)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

(1)

Page: 108
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Health Indicators (2, 3)											
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Gestational Diabetes During Pregnancy	Hypertension During Pregnancy	
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester	Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1955	160	7.5 (169)	53.1 (81)	128	31.3 (135)	39.1 (21)	17	* *	44	* *	59	* * 59
1957	518	5.4 (141)	53.1 (81)	451	18.2 (13)	55.0 (155)	67	* *	132	13.6 (3)	199	5.5 (47) 199
1959	224	4.0 (61)	58.0 (144)	191	20.4 (43)	50.8 (113)	39	* *	50	* *	67	* * 67
1963	35	* * * *	* *	24	* * * *	* *	3	* *	6	* *	0	* * 0
2150	802	3.0 (27)	54.1 (100)	713	25.9 (102)	48.1 (78)	76	* *	193	16.6 (7)	204	12.7 (114) 204
2151	334	5.7 (152)	48.5 (19)	290	25.2 (94)	52.1 (126)	40	* *	75	* *	78	* * 78
2251	539	3.0 (27)	47.9 (15)	467	29.8 (130)	43.7 (41)	53	* *	57	* *	121	3.3 (5) 121
2252	274	4.0 (61)	52.9 (76)	230	69.1 (169)	17.4 (3)	12	* *	50	* *	62	* * 62
2551	4,353	4.8 (106)	53.5 (91)	3,015	21.5 (61)	51.9 (124)	840	54.0 (70)	1,103	43.2 (84)	476	5.3 (40) 476
2552	8	* * * *	* *	0	* * * *	* *	1	* *	0	* *	0	* * 0
2554	135	8.1 (171)	53.3 (88)	112	57.1 (163)	21.4 (5)	11	* *	19	* *	26	* * 27
2555	4,791	5.2 (130)	53.3 (88)	3,500	19.4 (30)	54.6 (152)	920	37.6 (45)	1,487	29.1 (46)	803	9.1 (102) 803
3001	1,128	4.0 (61)	57.7 (141)	954	25.4 (97)	46.0 (54)	148	39.2 (50)	623	31.5 (55)	303	3.0 (1) 305
3002	266	3.0 (27)	61.3 (166)	217	28.1 (118)	47.9 (75)	23	* *	157	43.3 (85)	72	* * 72
3003	6	* * * *	* *	4	* * * *	* *	2	* *	1	* *	4	* * 4
3004	217	0.5 (1)	65.9 (171)	183	22.4 (68)	46.4 (63)	17	* *	38	* *	45	* * 45
3007	36	* * * *	* *	28	* * * *	* *	4	* *	15	* *	7	* * 7

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Page: 109
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Health Indicators (2, 3)											
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Gestational Diabetes During Pregnancy	Hypertension During Pregnancy	
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester	Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
3009	14	* *	* *	10	* *	* *	1	* *	4	* *	3	* *
3010	2	* *	* *	0	* *	* *	0	* *	0	* *	0	* *
3201	671	3.9 (54)	53.8 (93)	596	21.6 (63)	54.4 (150)	56	* *	96	* *	182	9.3 (103)
3301	4,752	4.4 (87)	52.0 (58)	3,913	33.5 (144)	44.6 (46)	946	30.4 (21)	1,254	38.3 (77)	2,372	4.9 (32)
3302	342	2.3 (8)	60.8 (163)	291	29.9 (131)	44.0 (43)	60	* *	85	* *	155	7.7 (87)
3303	14	* *	* *	14	* *	* *	2	* *	5	* *	0	* *
3304	123	9.8 (173)	35.8 (2)	109	41.3 (158)	31.2 (9)	19	* *	41	* *	51	* *
3305	120	4.2 (77)	39.2 (3)	108	35.2 (147)	35.2 (16)	16	* *	46	* *	57	* *
3306	201	4.5 (96)	60.7 (162)	177	24.9 (89)	51.4 (119)	32	* *	65	* *	113	7.1 (76)
3308	60	* *	* *	44	* *	* *	12	* *	19	* *	20	* *
3309	42	* *	* *	41	* *	* *	12	* *	14	* *	18	* *
3311	12	* *	* *	11	* *	* *	1	* *	2	* *	0	* *
3550	92	* *	* *	80	* *	* *	10	* *	17	* *	23	* *
3551	35	* *	* *	30	* *	* *	2	* *	9	* *	6	* *
3552	53	* *	* *	51	* *	* *	3	* *	18	* *	11	* *
3701	957	4.5 (96)	53.1 (81)	797	18.3 (16)	55.7 (160)	145	22.8 (8)	173	24.9 (26)	420	6.2 (64)
3702	126	4.8 (106)	59.5 (152)	110	20.9 (53)	53.6 (141)	25	* *	21	* *	0	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Page: 110
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Health Indicators (2, 3)											
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Gestational Diabetes During Pregnancy	Hypertension During Pregnancy	
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester	Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
3703	54	* *	* *	48	* *	* *	9	* *	11	* *	0	* *
3704	51	* *	* *	46	* *	* *	10	* *	8	* *	0	* *
3805	3,960	4.9 (116)	51.0 (47)	3,524	18.6 (20)	53.7 (143)	514	33.1 (28)	669	36.9 (72)	1,868	4.2 (15)
3950	2,443	4.2 (77)	54.8 (111)	2,094	18.3 (16)	54.1 (146)	352	37.2 (43)	547	32.9 (61)	878	8.0 (92)
3952	47	* *	* *	44	* *	* *	4	* *	12	* *	4	* *
3953	36	* *	* *	31	* *	* *	3	* *	11	* *	5	* *
3954	131	5.3 (133)	52.7 (72)	120	25.8 (100)	50.0 (93)	19	* *	24	* *	26	* *
4001	1,237	4.2 (77)	58.3 (146)	987	34.7 (146)	42.4 (33)	145	44.8 (61)	164	40.2 (80)	465	4.7 (27)
4002	1,068	3.2 (34)	53.2 (86)	902	26.5 (106)	49.9 (92)	134	48.5 (64)	146	45.2 (90)	400	10.0 (107)
4103	123	1.6 (3)	69.9 (172)	105	29.5 (129)	41.0 (26)	17	* *	16	* *	46	* *
4106	1,154	3.2 (34)	57.4 (139)	992	24.9 (89)	47.4 (71)	145	47.6 (63)	209	39.7 (79)	360	5.8 (53)
4107	3,880	3.8 (52)	50.2 (31)	3,212	20.7 (49)	51.5 (121)	574	43.6 (59)	929	30.2 (50)	1,686	5.2 (38)
4109	1,272	4.5 (96)	49.8 (27)	940	24.8 (86)	48.8 (85)	194	23.7 (11)	314	16.6 (7)	536	6.2 (64)
4110	1,953	2.9 (24)	55.2 (115)	1,629	25.1 (92)	44.9 (50)	224	34.8 (35)	456	32.2 (57)	767	4.0 (10)
4111	1	* *	* *	0	* *	* *	0	* *	0	* *	0	* *
4112	703	1.6 (3)	60.5 (159)	640	62.8 (166)	16.3 (2)	56	* *	132	72.7(100)	283	10.6 (110)
4113	950	2.4 (12)	56.6 (131)	856	26.6 (107)	41.6 (29)	69	* *	156	44.9 (88)	302	6.3 (67)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Page: 111
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Health Indicators (2, 3)											
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Gestational Diabetes During Pregnancy	Hypertension During Pregnancy	
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester	Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4120	2,340	5.3 (133)	50.6 (42)	1,892	18.9 (25)	51.1 (118)	327	30.0 (19)	590	21.4 (14)	914	4.3 (17)
4160	2,057	4.5 (96)	49.3 (22)	1,605	24.5 (82)	47.2 (70)	345	30.4 (21)	617	25.4 (28)	811	6.9 (74)
4351	752	3.9 (54)	54.0 (96)	663	20.2 (42)	53.7 (143)	78	* *	194	25.8 (30)	181	4.4 (22)
4352	212	2.4 (12)	55.7 (120)	183	25.7 (99)	50.3 (102)	23	* *	61	* *	52	* *
4353	586	3.6 (45)	53.8 (93)	504	26.0 (103)	50.2 (99)	71	* *	149	24.8 (25)	147	7.5 (83)
4354	927	3.7 (49)	53.0 (78)	804	20.6 (46)	53.6 (141)	106	30.2 (20)	232	39.2 (78)	245	5.3 (40)
4355	411	5.1 (122)	54.0 (96)	354	14.4 (1)	61.6 (169)	51	* *	84	* *	97	* *
4356	569	5.1 (122)	56.6 (131)	495	16.2 (4)	58.6 (168)	59	* *	118	37.3 (73)	159	5.7 (51)
4357	13	* *	* *	12	* *	* *	0	* *	2	* *	2	* *
4358	228	1.3 (2)	71.5 (173)	165	39.4 (153)	32.1 (10)	18	* *	41	* *	63	* *
4359	615	4.4 (87)	56.7 (133)	533	27.6 (111)	46.3 (62)	65	* *	117	44.4 (87)	160	7.5 (83)
4360	208	2.4 (12)	60.1 (157)	178	20.8 (51)	48.9 (86)	19	* *	55	* *	44	* *
4361	298	4.7 (103)	51.3 (49)	259	20.1 (39)	57.5 (165)	36	* *	75	* *	75	* *
4362	395	4.1 (70)	50.9 (46)	357	19.3 (28)	55.7 (160)	46	* *	124	25.8 (30)	106	4.7 (27)
4363	1,100	3.9 (54)	49.7 (25)	981	19.6 (33)	54.3 (148)	98	* *	297	24.2 (23)	263	6.5 (70)
4401	1,197	6.2 (162)	51.6 (56)	1,058	18.8 (23)	52.6 (132)	155	32.9 (27)	340	32.4 (58)	409	5.9 (57)
4501	1,090	4.0 (61)	55.5 (118)	978	24.1 (77)	48.4 (79)	79	* *	204	16.7 (9)	309	7.8 (90)
											309	8.7 (43)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

Table 5A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Page: 112
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Health Indicators (2, 3)											
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Gestational Diabetes During Pregnancy	Hypertension During Pregnancy	
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester	Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4601	1,669	4.8 (106)	53.0 (78)	1,458	24.8 (86)	44.6 (46)	237	21.5 (7)	411	27.3 (37)	614	8.3 (97)
4701	1,469	4.8 (106)	47.3 (12)	1,312	17.1 (9)	58.5 (166)	233	13.7 (2)	497	16.1 (4)	502	6.0 (59)
5050	1,464	5.9 (157)	50.1 (30)	452	24.3 (81)	46.0 (54)	341	36.7 (41)	267	27.3 (37)	173	9.8 (105)
5051	3,070	5.1 (122)	50.3 (34)	1,086	22.7 (69)	50.4 (104)	624	38.0 (47)	590	28.8 (44)	400	6.8 (73)
5052	2,193	4.4 (87)	51.1 (48)	640	24.1 (77)	50.9 (114)	479	37.8 (46)	384	23.7 (20)	261	7.7 (87)
5054	2	* * * *	*	0	* * * *	*	0	*	0	* *	0	* *
5101	1,832	4.0 (61)	47.9 (15)	1,581	18.8 (23)	50.7 (111)	289	36.0 (40)	940	28.8 (44)	808	7.4 (81)
5202	656	2.4 (12)	50.2 (31)	587	19.4 (30)	50.1 (97)	83	*	159	20.8 (13)	184	8.2 (95)
5203	212	3.3 (40)	63.7 (170)	188	18.6 (20)	54.3 (148)	18	*	49	* *	42	* *
5204	198	5.1 (122)	52.5 (69)	159	15.1 (2)	50.9 (114)	10	*	45	* *	47	* *
5205	34	* * * *	*	28	* * * *	*	6	*	4	* *	0	* *
5601	1,332	3.8 (52)	52.3 (64)	1,181	19.3 (28)	52.8 (134)	152	13.2 (1)	360	47.8 (93)	382	7.1 (76)
5801	2,439	5.4 (141)	50.5 (38)	2,071	21.8 (66)	51.8 (122)	325	28.9 (17)	489	26.8 (36)	869	6.4 (68)
5803	22	* * * *	*	21	* * * *	*	0	*	12	* *	17	* *
5804	9	* * * *	*	9	* * * *	*	0	*	2	* *	7	* *
6101	44	* * * *	*	38	* * * *	*	8	*	4	* *	17	* *
6102	138	6.5 (165)	56.5 (129)	126	20.6 (46)	55.6 (159)	22	*	21	* *	67	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Page: 113
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Health Indicators (2, 3)											
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Gestational Diabetes During Pregnancy	Hypertension During Pregnancy	
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester	Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
6103	4,602	4.1 (70)	54.7 (110)	3,974	18.2 (13)	53.2 (139)	571	38.9 (49)	366	36.6 (70)	1,726	6.9 (74)
6104	71	* * * *	* *	55	* * * *	* *	11	* *	7	* *	39	* *
6105	38	* * * *	* *	36	* * * *	* *	5	* *	6	* *	19	* *
6106	2	* * * *	* *	2	* * * *	* *	0	* *	0	* *	4	* *
6301	5,500	4.1 (70)	53.2 (86)	4,860	19.6 (33)	50.6 (109)	942	33.1 (28)	1,547	23.6 (19)	1,703	6.2 (64)
6312	3,040	3.9 (54)	56.7 (133)	2,702	18.4 (18)	52.4 (130)	583	33.8 (32)	1,089	28.7 (42)	928	7.5 (83)
6316	1,554	5.0 (118)	50.7 (45)	1,396	16.4 (5)	54.0 (145)	254	22.8 (8)	612	20.3 (10)	509	4.5 (23)
6320	276	4.7 (103)	48.9 (21)	241	24.9 (89)	46.1 (56)	54	* *	83	* *	79	* *
6325	172	2.3 (8)	50.0 (28)	136	23.5 (71)	50.0 (93)	26	* *	58	* *	48	* *
6330	506	2.8 (23)	55.9 (121)	405	23.7 (72)	47.7 (73)	89	* *	106	33.0 (63)	174	5.7 (51)
6335	661	5.0 (118)	47.8 (14)	572	21.3 (58)	50.0 (93)	114	33.3 (30)	144	22.9 (17)	211	4.3 (17)
6340	210	4.3 (83)	61.0 (165)	159	23.9 (75)	43.4 (39)	30	* *	42	* *	49	* *
6345	58	* * * *	* *	49	* * * *	* *	11	* *	20	* *	19	* *
6350	305	3.9 (54)	47.2 (10)	237	24.1 (77)	51.9 (124)	57	* *	58	* *	92	* *
7304	246	4.5 (96)	59.3 (151)	212	21.2 (56)	50.9 (114)	25	* *	69	* *	71	* *
7306	2,889	4.8 (106)	57.1 (138)	2,408	20.8 (51)	51.4 (119)	433	42.0 (56)	555	45.2 (90)	789	7.1 (76)
7308	212	2.4 (12)	62.3 (167)	194	20.6 (46)	48.5 (83)	32	* *	49	* *	66	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

(1)

Page: 114
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Health Indicators (2, 3)											
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Gestational Diabetes During Pregnancy	Hypertension During Pregnancy	
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester	Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
7310	27	* * * *	* *	25	* * * *	* *	4	* *	4	* *	19	* *
7401	2,364	5.3 (133)	50.2 (31)	1,827	20.1 (39)	52.1 (126)	328	34.5 (33)	1,238	29.5 (49)	725	11.0 (111)
7402	280	4.3 (83)	49.6 (24)	213	16.9 (7)	47.9 (75)	54	* *	141	16.3 (5)	48	* *
7403	101	5.9 (157)	51.5 (53)	78	* * * *	* *	18	* *	54	* *	21	* *
7404	103	5.8 (155)	46.6 (9)	85	* * * *	* *	11	* *	72	* *	17	* *
7601	837	5.5 (146)	52.9 (76)	724	20.9 (53)	49.0 (87)	91	* *	197	23.9 (22)	230	10.0 (107)
7850	763	5.8 (155)	52.2 (62)	669	18.1 (11)	54.4 (150)	114	31.6 (26)	179	28.5 (41)	369	4.3 (17)
7851	165	7.3 (168)	44.8 (8)	147	17.7 (10)	53.1 (137)	30	* *	32	* *	56	* *
7852	103	3.9 (54)	56.3 (124)	92	* * * *	* *	16	* *	29	* *	38	* *
8150	69	* * * *	* *	56	* * * *	* *	12	* *	15	* *	20	* *
8151	3,144	4.4 (87)	54.5 (106)	2,582	24.1 (77)	47.8 (74)	520	39.6 (53)	822	31.8 (56)	1,108	7.8 (90)
8152	568	7.2 (167)	42.8 (4)	465	34.2 (145)	42.6 (34)	63	* *	73	* *	189	3.2 (4)
8153	36	* * * *	* *	31	* * * *	* *	11	* *	14	* *	8	* *
8154	351	8.8 (172)	31.9 (1)	293	25.3 (96)	39.2 (22)	55	* *	108	28.7 (42)	91	* *
8155	96	* * * *	* *	88	* * * *	* *	12	* *	25	* *	21	* *
8156	46	* * * *	* *	40	* * * *	* *	6	* *	12	* *	10	* *
8157	25	* * * *	* *	18	* * * *	* *	5	* *	4	* *	10	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

(1)

Page: 115
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Health Indicators (2, 3)											
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Gestational Diabetes During Pregnancy	Hypertension During Pregnancy	
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester	Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8201	2,535	5.2 (130)	52.1 (60)	1,408	24.7 (84)	50.6 (109)	633	24.3 (13)	557	23.7 (20)	603	8.0 (92)
8202	3,355	4.0 (61)	47.2 (10)	2,187	39.8 (154)	34.0 (13)	571	31.2 (24)	575	27.5 (39)	853	5.3 (40)
8203	304	2.6 (20)	54.9 (112)	163	31.9 (137)	46.6 (64)	55	* *	56	* *	53	* * 53
8250	332	4.5 (96)	49.4 (23)	207	15.9 (3)	50.2 (99)	99	* *	74	* *	218	6.4 (68)
8255	3,017	5.4 (141)	52.3 (64)	1,492	37.3 (151)	40.9 (25)	704	25.7 (14)	646	26.0 (32)	328	6.1 (60)
8257	553	6.1 (161)	44.5 (7)	317	42.3 (160)	32.2 (11)	104	44.2 (60)	106	26.4 (34)	136	6.6 (72)
8258	1,736	2.9 (24)	54.5 (106)	864	40.4 (157)	38.9 (19)	365	34.5 (33)	214	26.6 (35)	326	7.4 (81)
8260	473	5.5 (146)	43.3 (6)	429	28.9 (127)	39.4 (23)	78	* *	110	38.2 (76)	161	12.4 (113)
8901	1,042	4.8 (106)	58.4 (149)	861	28.3 (122)	49.2 (89)	117	61.5 (71)	92	* *	297	3.0 (1)
8902	601	2.7 (22)	60.1 (157)	502	27.3 (109)	46.6 (64)	99	* *	91	* *	213	8.9 (101)
8903	1,202	3.2 (34)	59.7 (155)	996	27.6 (111)	46.2 (60)	169	52.7 (69)	109	51.4 (96)	398	5.8 (53)
8904	927	2.5 (19)	59.5 (152)	795	32.2 (138)	40.8 (24)	118	41.5 (55)	127	53.5 (97)	289	10.4 (109)
8905	2,451	3.7 (49)	54.1 (100)	2,072	28.3 (122)	43.1 (37)	109	45.9 (62)	154	46.1 (92)	907	4.3 (17)
9004	317	5.0 (118)	56.8 (136)	284	20.4 (43)	54.9 (154)	63	* *	73	* *	3	* * 3
9010	35	* *	* *	31	* *	* *	2	* *	4	* *	0	* * 0
9012	14	* *	* *	11	* *	* *	0	* *	0	* *	13	* * 13
9013	30	* *	* *	26	* *	* *	4	* *	3	* *	32	* * 32

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

(1)

Page: 116
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Health Indicators (2, 3)											
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Gestational Diabetes During Pregnancy	Hypertension During Pregnancy	
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester	Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
9015	8	* *	* *	7	* *	* *	2	* *	0	* *	10	* * 10
9016	35	* *	* *	27	* *	* *	4	* *	7	* *	30	* * 30
9017	8	* *	* *	6	* *	* *	1	* *	1	* *	10	* * 10
9018	31	* *	* *	28	* *	* *	0	* *	1	* *	31	* * 31
9019	12	* *	* *	10	* *	* *	1	* *	0	* *	12	* * 12
9021	14	* *	* *	13	* *	* *	3	* *	0	* *	20	* * 20
9023	80	* *	* *	69	* *	* *	7	* *	0	* *	0	* * 0
9040	268	2.2 (6)	60.8 (163)	240	25.8 (100)	52.5 (131)	54	* *	38	* *	6	* * 6
9041	842	3.6 (45)	63.3 (169)	639	35.4 (148)	43.2 (38)	41	* *	66	* *	143	5.6 (48) 143
9045	1,344	4.8 (106)	50.5 (38)	1,153	32.3 (139)	42.8 (36)	107	35.5 (37)	127	36.2 (69)	438	3.4 (7) 439
9050	2,188	3.3 (40)	58.3 (146)	1,908	27.8 (114)	45.3 (52)	273	38.5 (48)	327	33.6 (65)	720	5.4 (45) 721
9051	2,981	3.2 (34)	54.9 (112)	2,500	28.7 (125)	42.6 (34)	164	28.7 (15)	362	33.4 (64)	827	7.5 (83) 825
9053	3,296	4.0 (61)	59.6 (154)	2,828	32.4 (140)	45.9 (53)	611	42.4 (58)	818	37.4 (74)	992	4.1 (13) 990
9054	81	* *	* *	77	* *	* *	11	* *	5	* *	0	* * 0
9055	99	* *	* *	94	* *	* *	11	* *	0	* *	21	* * 22
9058	611	4.4 (87)	47.5 (13)	507	37.5 (152)	33.7 (12)	72	* *	90	* *	228	15.4 (117) 228
9066	2,772	4.3 (83)	58.0 (144)	2,476	32.6 (141)	43.5 (40)	388	42.0 (56)	553	49.0 (95)	856	5.0 (35) 861
												15.0 (114)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 5A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Maternal Health Indicators by Clinic
Combined 3 Years

Page: 117
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Health Indicators (2, 3)											
	Prepregnancy BMI (4)			Weight Gain (5)			Anemia (Low Hb/Hct) (6)			Gestational Diabetes During Pregnancy	Hypertension During Pregnancy	
	Under-weight	Over-weight	< Ideal	> Ideal	3rd Trimester	Postpartum						
	Number	% (Rank)	% (Rank)	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
9068	1,641	4.1 (70)	55.5 (118)	1,460	27.9 (116)	46.6 (64)	225	52.4 (68)	428	55.4 (98)	533	4.9 (32)
9069	2,563	3.1 (31)	60.6 (161)	2,249	22.3 (67)	52.2 (129)	456	34.9 (36)	729	32.6 (60)	1,032	4.8 (31)
9070	2,370	6.0 (160)	54.4 (105)	2,028	30.0 (132)	44.1 (44)	276	40.9 (54)	492	37.4 (74)	761	4.7 (27)
9072	1,933	3.1 (31)	57.7 (141)	1,771	25.2 (94)	49.5 (91)	281	39.5 (51)	426	45.1 (89)	682	5.3 (40)
9074	1,834	3.7 (49)	60.0 (156)	1,615	21.7 (65)	52.6 (132)	383	48.6 (65)	533	42.0 (81)	644	4.7 (27)
9079	526	5.1 (122)	57.0 (137)	431	26.0 (103)	47.1 (69)	117	23.9 (12)	82	* *	0	*
9080	2,530	4.2 (77)	57.5 (140)	2,158	21.4 (60)	50.5 (107)	362	48.9 (66)	414	42.8 (83)	805	6.1 (60)
Unknown Clinics	31	*	*	21	*	*	5	*	24	*	20	*
Michigan	186,664	4.3	53.6	150,285	25.4	48.1	27,292	34.5	42,450	32.0	59,987	6.2
											60,028	10.2

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight (BMI <18.5), normal weight (BMI 18.5 to 24.9), overweight (BMI 25.0 to 29.9), obese (BMI >30.0).

(5) Based on 2009 IOM report, "Weight Gain During Pregnancy": underweight during pregnancy (ideal weight gain = 28 to 40 pounds), normal weight prepregnancy (ideal weight gain = 25 to 35 pounds), overweight prepregnancy (ideal weight gain = 15 to 25 pounds), obese prepregnancy (ideal weight gain = 11 to 20 pounds).

(6) Based on 1990 MMWR, "Recommendations to Prevent and Control Iron Deficiency in the United States"; adjusted for altitude and smoking.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

(1)

Page: 118
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Behavioral Indicators (2, 3)					
	Medical Care		WIC Enrollment		Multivitamin Consumption	
	1st Trimester	No Care	1st Trimester		Prepregnancy (4)	During Pregnancy (5)
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number % (Rank)
0101	141	90.8 (13)	2.8 (43)	158	45.6 (26)	9 * * 11 * *
0102	445	92.1 (3)	1.1 (5)	491	42.8 (51)	40 * * 47 * *
0103	465	91.8 (6)	2.6 (38)	501	43.9 (39)	39 * * 40 * *
0104	133	91.7 (8)	1.5 (9)	143	45.5 (28)	13 * * 14 * *
0250	169	85.2 (57)	5.3 (92)	182	58.8 (4)	11 * * 14 * *
0251	148	87.2 (39)	8.8 (140)	159	57.9 (5)	13 * * 16 * *
0252	139	79.9 (119)	8.6 (137)	146	57.5 (6)	6 * * 9 * *
0253	185	89.2 (23)	2.7 (40)	197	58.9 (3)	13 * * 16 * *
0401	596	89.3 (21)	2.5 (34)	631	37.2 (93)	63 * * 66 * *
0402	498	92.2 (2)	2.8 (43)	528	45.3 (30)	38 * * 42 * *
0403	134	88.1 (30)	4.5 (78)	146	51.4 (10)	6 * * 6 * *
0404	155	90.3 (15)	3.9 (64)	162	42.0 (61)	10 * * 12 * *
0501	572	85.8 (50)	8.2 (133)	624	49.8 (13)	45 * * 51 * *
0502	4 * * * *			4 * * *	0 * * 0 * *	
0503	177	81.9 (101)	10.7 (154)	189	42.9 (50)	17 * * 17 * *
0504	604	89.6 (18)	4.3 (71)	660	47.9 (15)	44 * * 47 * *
0505	393	83.2 (84)	10.2 (152)	436	46.1 (23)	38 * * 40 * *
0506	198	73.2 (151)	15.7 (166)	236	56.4 (7)	23 * * 24 * *
0650	1,165	86.5 (44)	3.6 (59)	1,276	38.2 (83)	242 21.1 (24) 248 89.1 (24)
0651	669	81.2 (108)	4.6 (83)	755	42.8 (51)	141 21.3 (23) 158 80.4 (72)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

(1)

Page: 119
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Behavioral Indicators ^(2, 3)								
	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester	No Care	1st Trimester	Prepregnancy (4)		During Pregnancy (5)			
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0652	351	84.6 (63)	6.3 (108)	390	42.1 (60)	60	* * *	70	* * *
0653	458	87.3 (38)	4.4 (74)	503	47.5 (17)	103	21.4 (22)	105	94.3 (3)
0654	419	84.0 (74)	6.4 (113)	447	52.1 (9)	95	* * *	98	* * *
0655	450	87.6 (33)	5.3 (92)	484	36.2 (99)	76	* * *	79	* * *
0657	77	* * * * *		80	* * *	18	* * *	18	* * *
0701	434	79.5 (122)	7.8 (126)	496	37.7 (89)	30	* * *	30	* * *
0702	100	91.0 (12)	6.0 (102)	110	47.3 (19)	12	* * *	12	* * *
0703	271	86.3 (45)	8.5 (136)	293	46.8 (22)	22	* * *	24	* * *
0704	76	* * * * *		86	* * *	5	* * *	5	* * *
0705	189	82.0 (98)	5.8 (98)	209	36.4 (97)	15	* * *	15	* * *
0850	1,223	85.4 (54)	4.3 (71)	1,359	34.1 (107)	396	17.7 (38)	402	84.8 (51)
0851	965	85.1 (59)	6.4 (113)	1,046	40.7 (71)	311	13.2 (60)	319	84.6 (52)
0901	1,938	86.7 (43)	3.7 (61)	2,102	41.0 (67)	171	22.8 (15)	259	91.5 (14)
0902	82	* * * * *		83	* * *	6	* * *	6	* * *
1051	1,458	77.9 (133)	13.2 (163)	1,712	45.2 (31)	137	24.8 (9)	150	85.3 (50)
1055	1	* * * * *		2	* * *	0	* * *	0	* * *
1101	1,925	75.9 (138)	5.0 (86)	2,237	32.0 (115)	506	10.7 (67)	510	88.8 (27)
1102	948	85.4 (54)	0.7 (3)	1,043	32.7 (114)	208	15.4 (52)	228	93.9 (7)
1106	135	83.7 (80)	1.5 (9)	152	24.3 (145)	26	* * *	26	* * *
1250	1,072	82.9 (87)	6.2 (106)	1,230	43.8 (40)	249	24.9 (8)	302	67.5 (85)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

(1)

Page: 120
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Behavioral Indicators (2, 3)							
	Medical Care			WIC Enrollment		Multivitamin Consumption		
	1st Trimester	No Care	1st Trimester	Prepregnancy (4)		During Pregnancy (5)		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number
1251	910	82.4 (95)	7.3 (124)	1,016	39.2 (79)	227	16.3 (48)	244
1252	800	78.4 (128)	8.9 (141)	897	42.0 (61)	207	21.7 (19)	211
1253	909	81.4 (106)	7.5 (125)	1,007	35.5 (103)	223	22.4 (16)	228
1350	2,697	82.0 (98)	8.0 (130)	3,006	39.3 (78)	879	29.4 (5)	911
1351	165	84.2 (69)	6.1 (104)	177	35.6 (102)	63	*	*
1352	482	77.2 (136)	11.0 (155)	520	43.7 (42)	160	16.9 (42)	160
1401	252	84.5 (65)	4.8 (84)	279	50.9 (11)	21	*	*
1402	162	75.3 (141)	11.1 (156)	193	45.1 (32)	17	*	*
1403	29	*	*	32	*	2	*	*
1404	14	*	*	25	*	3	*	*
1501	1,148	81.2 (108)	5.9 (99)	1,389	40.2 (73)	281	17.1 (40)	309
1502	10	*	*	14	*	0	*	*
1503	1,231	77.6 (135)	5.2 (90)	1,632	37.7 (89)	217	35.0 (2)	367
1504	2,734	89.3 (21)	1.3 (7)	3,000	37.3 (92)	688	15.3 (55)	722
1505	911	90.1 (16)	3.0 (51)	979	35.8 (101)	255	29.4 (5)	258
1506	661	88.7 (26)	2.9 (48)	738	37.4 (91)	184	10.9 (66)	187
1507	730	92.1 (3)	2.2 (27)	788	37.1 (94)	175	21.7 (19)	180
1508	142	88.7 (26)	1.4 (8)	241	31.1 (118)	32	*	*
1509	629	83.8 (79)	4.1 (68)	825	44.1 (37)	163	19.0 (34)	177
1510	203	85.2 (57)	0.5 (2)	234	42.3 (54)	37	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

(1)

Page: 121
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Behavioral Indicators (2, 3)									
	Medical Care			WIC Enrollment			Multivitamin Consumption			
	1st Trimester	No Care	1st Trimester	Prepregnancy (4)		During Pregnancy (5)				
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
1511	669	88.9 (25)	1.2 (6)	721	40.2 (73)	194	20.1 (28)	194	85.6 (46)	
1512	310	91.3 (11)	1.9 (22)	332	34.3 (106)	68	*	68	*	*
1701	670	78.2 (131)	11.5 (158)	762	47.2 (20)	26	*	44	*	*
1712	81	*	*	94	*	5	*	6	*	*
1950	738	82.8 (88)	9.3 (146)	817	41.6 (64)	174	19.0 (34)	178	92.7 (9)	
1952	176	84.1 (72)	4.0 (66)	190	27.4 (131)	23	*	25	*	*
1953	372	86.0 (48)	6.2 (106)	396	37.9 (86)	65	*	65	*	*
1954	711	84.5 (65)	5.5 (94)	758	42.2 (58)	157	18.5 (36)	179	83.2 (61)	
1955	157	83.4 (83)	9.6 (147)	171	47.4 (18)	31	*	34	*	*
1957	488	84.2 (69)	5.5 (94)	530	38.1 (84)	101	24.8 (9)	103	85.4 (47)	
1959	212	82.5 (93)	6.6 (118)	227	31.3 (117)	38	*	41	*	*
1963	29	*	*	36	*	0	*	0	*	*
2150	752	89.0 (24)	4.4 (74)	816	44.6 (33)	51	*	55	*	*
2151	297	79.5 (122)	11.8 (159)	351	42.2 (58)	25	*	31	*	*
2251	510	84.9 (60)	6.1 (104)	557	47.9 (15)	25	*	35	*	*
2252	256	91.8 (6)	1.6 (13)	278	59.0 (2)	16	*	22	*	*
2551	3,322	73.9 (148)	4.0 (66)	4,449	23.7 (147)	400	6.3 (80)	519	84.4 (54)	
2552	0	*	*	8	*	0	*	0	*	*
2554	122	82.8 (88)	4.1 (68)	135	43.0 (49)	13	*	13	*	*
2555	3,971	82.0 (98)	3.7 (61)	4,982	24.5 (142)	507	12.0 (64)	617	85.4 (47)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

(1)

Page: 122
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Behavioral Indicators (2, 3)								
	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester	No Care	1st Trimester			Prepregnancy (4)	During Pregnancy (5)		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
3001	1,007	71.1 (157)	9.7 (149)	1,172	46.0 (25)	119	6.7 (78)	121	83.5 (58)
3002	254	82.7 (91)	2.8 (43)	275	50.5 (12)	32	* *	32	* *
3003	7	*	*	10	*	1	*	1	*
3004	181	79.6 (121)	14.4 (165)	225	59.1 (1)	24	*	24	*
3007	32	*	*	41	*	2	*	2	*
3009	11	*	*	19	*	0	*	0	*
3010	1	*	*	3	*	0	*	0	*
3201	640	92.0 (5)	1.7 (17)	677	45.5 (28)	76	*	78	*
3301	4,606	83.0 (86)	2.9 (48)	5,002	25.7 (137)	1,429	16.4 (47)	1,621	86.8 (40)
3302	340	80.3 (117)	2.9 (48)	355	21.1 (156)	51	*	99	*
3303	15	*	*	15	*	0	*	0	*
3304	124	85.5 (52)	3.2 (54)	127	26.0 (134)	31	*	32	*
3305	122	86.9 (40)	1.6 (13)	127	26.0 (134)	25	*	27	*
3306	198	89.4 (20)	2.5 (34)	208	20.7 (157)	57	*	60	*
3308	55	*	*	62	*	13	*	14	*
3309	41	*	*	42	*	11	*	11	*
3311	12	*	*	11	*	0	*	0	*
3550	83	*	*	91	*	10	*	11	*
3551	34	*	*	37	*	2	*	2	*
3552	54	*	*	52	*	0	*	0	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

(1)

Page: 123
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Behavioral Indicators (2, 3)									
	Medical Care			WIC Enrollment			Multivitamin Consumption			
	1st Trimester	No Care	1st Trimester	Prepregnancy (4)		During Pregnancy (5)				
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
3701	1,024	87.4 (37)	2.2 (27)	1,099	36.8 (95)	162	11.7 (65)	232	91.4 (15)	
3702	118	84.7 (62)	6.8 (121)	126	43.7 (42)	0	*	*	0	*
3703	51	*	*	53	*	0	*	*	0	*
3704	55	*	*	51	*	0	*	*	0	*
3805	3,731	87.5 (35)	2.4 (33)	3,991	38.7 (82)	1,171	16.2 (49)	1,266	77.6 (79)	
3950	2,226	83.6 (81)	5.0 (86)	2,521	37.8 (87)	530	13.0 (62)	535	84.5 (53)	
3952	49	*	*	50	*	2	*	*	2	*
3953	35	*	*	40	*	1	*	*	1	*
3954	127	88.2 (29)	7.1 (122)	134	43.3 (45)	15	*	*	15	*
4001	1,141	80.5 (115)	3.3 (56)	1,302	44.5 (35)	280	12.9 (63)	285	67.0 (86)	
4002	989	84.0 (74)	2.3 (29)	1,106	40.1 (75)	233	20.6 (27)	241	77.2 (81)	
4103	116	66.4 (167)	12.9 (162)	124	33.1 (112)	15	*	*	15	*
4106	1,108	73.5 (150)	12.5 (161)	1,203	34.1 (107)	167	9.6 (71)	183	86.3 (42)	
4107	3,736	81.3 (107)	8.0 (130)	4,043	36.4 (97)	848	22.1 (18)	1,014	91.0 (18)	
4109	1,301	82.2 (96)	9.1 (144)	1,409	35.5 (103)	288	24.0 (11)	296	88.2 (31)	
4110	1,911	75.7 (140)	12.3 (160)	2,083	40.8 (68)	386	15.5 (51)	473	87.1 (39)	
4111	0	*	*	1	*	0	*	*	0	*
4112	731	54.4 (171)	4.4 (74)	804	28.7 (123)	155	25.8 (7)	181	55.2 (88)	
4113	857	50.9 (172)	29.3 (171)	987	46.1 (23)	149	3.4 (82)	167	94.0 (5)	
4120	2,306	81.1 (110)	9.0 (143)	2,446	33.5 (110)	524	16.8 (43)	536	91.2 (16)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

(1)

Page: 124
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Behavioral Indicators (2, 3)								
	Medical Care			WIC Enrollment		Multivitamin Consumption			
	1st Trimester	No Care	1st Trimester			Prepregnancy (4)	During Pregnancy (5)		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4160	2,125	82.8 (88)	5.9 (99)	2,266	28.2 (128)	484	21.5 (21)	501	90.0 (20)
4351	711	88.0 (31)	2.1 (25)	772	42.4 (53)	68	* *	71	* *
4352	197	86.8 (41)	1.5 (9)	222	41.4 (66)	19	* *	21	* *
4353	540	82.6 (92)	7.2 (123)	631	43.7 (42)	52	* *	57	* *
4354	848	91.6 (10)	1.8 (19)	923	40.8 (68)	90	* *	101	89.1 (24)
4355	390	75.1 (145)	10.0 (150)	462	39.0 (80)	41	* *	44	* *
4356	596	84.1 (72)	8.2 (133)	666	44.6 (33)	68	* *	69	* *
4357	14	*	*	19	*	0	* *	0	* *
4358	193	71.0 (158)	21.2 (169)	274	52.6 (8)	24	* *	35	* *
4359	581	86.1 (47)	3.4 (57)	641	39.0 (80)	56	* *	56	* *
4360	190	80.5 (115)	7.9 (128)	211	44.5 (35)	13	* *	17	* *
4361	271	84.9 (60)	2.6 (38)	304	37.8 (87)	29	* *	30	* *
4362	371	85.4 (54)	6.7 (119)	405	41.5 (65)	29	* *	30	* *
4363	1,025	81.0 (112)	10.0 (150)	1,113	42.3 (54)	97	* *	126	86.5 (41)
4401	1,164	79.8 (120)	10.2 (152)	1,560	31.4 (116)	135	23.0 (13)	146	83.6 (57)
4501	1,112	86.0 (48)	4.1 (68)	1,145	43.2 (46)	114	21.1 (24)	117	95.7 (1)
4601	1,696	87.7 (32)	2.5 (34)	1,878	24.1 (146)	254	15.4 (52)	328	89.3 (23)
4701	1,400	83.9 (77)	6.5 (115)	1,489	28.6 (126)	234	20.1 (28)	236	88.1 (34)
5050	834	87.6 (33)	1.7 (17)	1,519	24.6 (141)	187	18.2 (37)	248	92.3 (10)
5051	1,878	86.8 (41)	2.8 (43)	3,168	28.7 (123)	348	20.1 (28)	481	88.4 (30)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

(1)

Page: 125
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Behavioral Indicators (2, 3)									
	Medical Care			WIC Enrollment			Multivitamin Consumption			
	1st Trimester	No Care	1st Trimester	Prepregnancy (4)		During Pregnancy (5)				
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
5052	1,207	83.5 (82)	1.9 (22)	2,251	25.4 (139)	315	16.8 (43)	427	89.0 (26)	
5054	0	*	*	2	*	0	*	0	*	*
5101	1,705	88.3 (28)	1.6 (13)	1,936	24.5 (142)	283	23.7 (12)	369	94.0 (5)	
5202	642	89.6 (18)	3.9 (64)	685	40.7 (71)	57	*	60	*	*
5203	209	81.8 (103)	9.6 (147)	237	45.6 (26)	13	*	13	*	*
5204	181	90.6 (14)	5.5 (94)	220	42.3 (54)	11	*	11	*	*
5205	32	*	*	39	*	0	*	0	*	*
5601	1,306	89.9 (17)	1.5 (9)	1,378	36.1 (100)	139	23.0 (13)	146	89.7 (21)	
5801	2,417	81.1 (110)	6.7 (119)	2,637	38.0 (85)	477	19.7 (31)	479	91.2 (16)	
5803	22	*	*	26	*	1	*	1	*	*
5804	11	*	*	10	*	0	*	0	*	*
6101	43	*	*	46	*	10	*	11	*	*
6102	155	83.2 (84)	6.5 (115)	164	28.0 (129)	45	*	45	*	*
6103	4,647	83.9 (77)	6.5 (115)	5,130	43.8 (40)	930	15.8 (50)	1,017	79.4 (77)	
6104	74	*	*	81	*	23	*	24	*	*
6105	42	*	*	44	*	13	*	13	*	*
6106	5	*	*	5	*	1	*	1	*	*
6301	5,106	79.2 (125)	4.9 (85)	5,612	23.3 (149)	675	17.0 (41)	815	84.4 (54)	
6312	2,882	81.6 (104)	3.1 (53)	3,105	18.9 (163)	354	19.5 (32)	424	90.8 (19)	
6316	1,488	84.3 (67)	3.8 (63)	1,559	21.2 (155)	189	22.2 (17)	232	87.5 (37)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

(1)

Page: 126
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Behavioral Indicators (2, 3)									
	Medical Care			WIC Enrollment			Multivitamin Consumption			
	1st Trimester	No Care	1st Trimester	Prepregnancy (4)		During Pregnancy (5)				
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
6320	276	86.2 (46)	5.1 (89)	303	24.4 (144)	8	*	*	36	*
6325	160	85.6 (51)	6.3 (108)	181	25.4 (139)	15	*	*	21	*
6330	460	75.9 (138)	4.3 (71)	523	28.7 (123)	76	*	*	88	*
6335	639	84.0 (74)	6.3 (108)	702	30.2 (121)	18	*	*	104	87.5 (37)
6340	182	84.6 (63)	6.0 (102)	219	32.9 (113)	4	*	*	17	*
6345	55	*	*	59	*	1	*	*	7	*
6350	288	78.8 (127)	6.3 (108)	329	23.1 (151)	17	*	*	45	*
7304	248	80.2 (118)	8.9 (141)	274	36.5 (96)	22	*	*	24	*
7306	2,855	78.3 (130)	9.1 (144)	3,186	39.7 (77)	360	19.4 (33)	375	82.9 (64)	
7308	218	91.7 (8)	5.0 (86)	229	33.2 (111)	30	*	*	32	*
7310	33	*	*	34	*	6	*	*	6	*
7401	2,103	75.2 (143)	14.3 (164)	2,457	41.8 (63)	341	15.2 (56)	375	79.7 (74)	
7402	269	82.5 (93)	7.8 (126)	287	31.0 (119)	21	*	*	22	*
7403	98	*	*	111	42.3 (54)	4	*	*	5	*
7404	98	*	*	110	35.5 (103)	11	*	*	13	*
7601	768	81.6 (104)	8.7 (139)	861	43.1 (47)	94	*	*	101	83.2 (61)
7850	1,022	62.0 (169)	31.1 (172)	1,156	47.1 (21)	195	15.4 (52)	195	86.2 (43)	
7851	193	69.4 (164)	24.4 (170)	211	43.1 (47)	39	*	*	39	*
7852	119	73.1 (153)	21.0 (168)	128	33.6 (109)	16	*	*	16	*
8150	73	*	*	82	*	9	*	*	9	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

(1)

Page: 127
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Behavioral Indicators (2, 3)									
	Medical Care			WIC Enrollment			Multivitamin Consumption			
	1st Trimester	No Care	1st Trimester	Prepregnancy (4)		During Pregnancy (5)				
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
8151	3,031	84.3 (67)	2.3 (29)	3,261	27.0 (133)	579	20.7 (26)	624	88.1 (34)	
8152	542	77.9 (133)	6.3 (108)	595	44.0 (38)	103	7.8 (75)	106	83.0 (63)	
8153	36	*	*	38	*	3	*	4	*	*
8154	323	92.9 (1)	2.5 (34)	360	29.7 (122)	38	*	39	*	*
8155	95	*	*	103	27.2 (132)	4	*	8	*	*
8156	43	*	*	50	*	5	*	5	*	*
8157	23	*	*	27	*	7	*	10	*	*
8201	2,007	84.2 (69)	1.9 (22)	2,611	19.8 (159)	475	15.2 (56)	494	91.9 (12)	
8202	2,825	79.2 (125)	8.6 (137)	3,511	40.8 (68)	339	16.5 (46)	542	81.2 (69)	
8203	216	87.5 (35)	2.3 (29)	316	15.8 (166)	61	*	74	*	*
8250	226	81.0 (112)	0.0 (1)	351	12.0 (173)	244	13.9 (59)	246	94.7 (2)	
8255	2,190	82.1 (97)	5.2 (90)	3,124	22.3 (153)	458	16.6 (45)	526	93.5 (8)	
8257	453	80.6 (114)	5.5 (94)	572	28.5 (127)	107	2.8 (83)	115	92.2 (11)	
8258	1,400	81.9 (101)	4.4 (74)	1,781	22.9 (152)	248	17.3 (39)	257	94.2 (4)	
8260	446	74.7 (147)	1.6 (13)	480	23.5 (148)	66	*	86	*	*
8901	1,006	68.3 (166)	4.5 (78)	1,133	19.9 (158)	114	9.6 (71)	149	88.6 (29)	
8902	629	76.2 (137)	3.2 (54)	699	19.6 (161)	42	*	96	*	*
8903	1,180	70.6 (160)	3.6 (59)	1,353	19.8 (159)	118	7.6 (77)	204	88.7 (28)	
8904	920	78.4 (128)	2.7 (40)	1,020	23.2 (150)	72	*	144	85.4 (47)	
8905	2,390	78.2 (131)	4.5 (78)	2,771	30.7 (120)	230	6.5 (79)	456	77.9 (78)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

(1)

Page: 128
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Behavioral Indicators (2, 3)											
	Medical Care			WIC Enrollment			Multivitamin Consumption					
	1st Trimester	No Care		1st Trimester			Prepregnancy (4)		During Pregnancy (5)			
	Number	% (Rank)	% (Rank)	Number	% (Rank)		Number	% (Rank)	Number	% (Rank)		
9004	308	66.2 (168)	8.4 (135)	329	12.5 (172)		0	*	*	3	*	*
9010	29	*	*	35	*	*	0	*	*	0	*	*
9012	15	*	*	14	*	*	9	*	*	10	*	*
9013	34	*	*	33	*	*	12	*	*	12	*	*
9015	11	*	*	11	*	*	8	*	*	9	*	*
9016	35	*	*	38	*	*	21	*	*	21	*	*
9017	10	*	*	11	*	*	6	*	*	8	*	*
9018	34	*	*	34	*	*	19	*	*	19	*	*
9019	13	*	*	14	*	*	6	*	*	10	*	*
9021	21	*	*	21	*	*	11	*	*	14	*	*
9023	66	*	*	80	*	*	0	*	*	0	*	*
9040	262	61.8 (170)	11.1 (156)	273	16.8 (164)		0	*	*	3	*	*
9041	850	79.4 (124)	1.8 (19)	976	19.2 (162)		37	*	*	42	*	*
9045	1,289	70.3 (161)	8.0 (130)	1,472	25.5 (138)		232	8.6 (74)		235	82.6 (65)	
9050	2,035	72.2 (156)	7.9 (128)	2,280	28.0 (129)		219	31.5 (3)		397	77.1 (82)	
9051	2,644	68.5 (165)	17.2 (167)	3,042	48.3 (14)		247	43.7 (1)		390	66.7 (87)	
9053	3,035	72.8 (154)	2.8 (43)	3,373	16.0 (165)		390	30.8 (4)		505	87.9 (36)	
9054	78	*	*	81	*	*	0	*	*	0	*	*
9055	105	75.2 (143)	1.0 (4)	113	13.3 (169)		1	*	*	2	*	*
9058	580	85.5 (52)	3.4 (57)	619	39.9 (76)		137	4.4 (81)		145	83.4 (59)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 6A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Maternal Behavioral Indicators by Clinic
Combined 3 Years

Page: 129
Run Date: 7 / 21 / 10

Clinic	Comparison of Maternal Behavioral Indicators ^(2, 3)							
	Medical Care			WIC Enrollment		Multivitamin Consumption		
	1st Trimester	No Care	1st Trimester	Prepregnancy (4)		During Pregnancy (5)		
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number
9066	2,646	73.2 (151)	1.8 (19)	2,842	13.3 (169)	220	10.5 (70)	444
9068	1,548	70.7 (159)	2.1 (25)	1,670	15.4 (167)	161	10.6 (68)	254
9069	2,496	73.6 (149)	3.0 (51)	2,660	10.4 (174)	428	13.1 (61)	556
9070	2,230	69.7 (163)	5.9 (99)	2,432	21.9 (154)	229	14.0 (58)	367
9072	1,876	72.4 (155)	2.3 (29)	1,973	10.4 (174)	334	7.8 (75)	374
9074	1,759	69.8 (162)	4.5 (78)	1,861	12.7 (171)	265	10.6 (68)	347
9079	493	75.3 (141)	4.5 (78)	530	14.3 (168)	0	*	*
9080	2,374	75.0 (146)	2.7 (40)	2,589	26.0 (134)	395	8.9 (73)	419
Unknown Clinics	34	*	*	459	8.5 (176)	8	*	*
Michigan	173,011	80.5	5.7	197,712	32.3	29,692	18.2	34,588
								85.2

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Women who reported taking a multivitamin four times a week or more in the month before pregnancy; proxy for folic acid consumption.

(5) Pregnant women who reported taking a multivitamin in the past month; proxy for prenatal iron consumption.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

(1)

Page: 130
Run Date: 7 / 21 / 10

Clinic	Comparison of Smoking Indicators (2, 3)									
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal	Postpartum			
Clinic	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0101	134	52.2 (145)	137	36.5 (160)	36	* *	10	* *	39	* *
0102	442	56.3 (158)	427	35.1 (159)	143	23.8 (91)	47	* *	124	7.3 (75)
0103	411	57.4 (162)	416	37.0 (162)	124	30.6 (64)	40	* *	123	8.9 (92)
0104	134	61.9 (169)	129	34.9 (158)	50	* *	14	* *	38	* *
0250	175	62.3 (170)	160	31.9 (144)	61	* *	12	* *	42	* *
0251	154	48.1 (114)	146	19.2 (58)	48	* *	15	* *	38	* *
0252	144	44.4 (90)	132	25.0 (91)	47	* *	9	* *	40	* *
0253	179	54.7 (152)	178	31.5 (142)	66	* *	14	* *	49	* *
0401	572	54.9 (153)	582	33.8 (154)	206	28.6 (75)	66	* *	155	4.5 (31)
0402	482	52.9 (147)	481	33.7 (152)	156	23.1 (95)	39	* *	136	4.4 (29)
0403	129	49.6 (125)	129	24.0 (85)	39	* *	6	* *	36	* *
0404	148	48.6 (120)	151	29.8 (132)	51	* *	12	* *	41	* *
0501	576	51.2 (136)	569	28.3 (119)	180	36.7 (35)	49	* *	145	12.4 (110)
0502	3	* *	4	* *	1	* *	0	* *	1	* *
0503	171	52.0 (141)	176	25.6 (96)	55	* *	17	* *	54	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

(1)

Page: 131
Run Date: 7 / 21 / 10

Clinic	Comparison of Smoking Indicators (2, 3)									
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal	Postpartum			
Clinic	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0504	608	50.0 (129)	601	33.1 (149)	190	39.5 (25)	45	* *	157	5.7 (51)
0505	383	49.3 (123)	395	30.4 (136)	124	28.2 (76)	39	* *	109	5.5 (49)
0506	226	52.2 (145)	205	29.3 (126)	73	* *	24	* *	53	* *
0650	1,005	51.9 (140)	1,010	27.6 (115)	307	30.0 (67)	247	19.0 (50)	465	7.1 (71)
0651	594	63.5 (171)	615	41.5 (169)	231	16.0 (102)	155	35.5 (87)	264	12.5 (111)
0652	276	65.6 (172)	297	39.1 (168)	112	25.9 (85)	70	* *	122	13.9 (114)
0653	457	49.9 (128)	454	27.3 (111)	149	28.9 (74)	105	30.5 (85)	176	20.5 (118)
0654	369	55.0 (154)	363	29.5 (128)	132	26.5 (84)	98	* *	163	14.7 (117)
0655	423	58.4 (165)	450	38.7 (167)	180	23.3 (94)	78	* *	153	9.2 (96)
0657	76	* *	71	* *	24	* *	18	* *	30	* *
0701	430	43.5 (86)	399	25.3 (94)	79	* *	30	* *	117	0.9 (1)
0702	104	67.3 (173)	97	* *	40	* *	10	* *	23	* *
0703	267	50.9 (134)	273	26.7 (104)	77	* *	24	* *	69	* *
0704	75	* *	75	* *	27	* *	5	* *	20	* *
0705	180	48.9 (121)	174	33.3 (150)	49	* *	15	* *	38	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

(1)

Page: 132
Run Date: 7 / 21 / 10

Clinic	Comparison of Smoking Indicators (2, 3)									
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
Clinic	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0850	937	57.1 (161)	1,026	27.7 (116)	325	24.6 (89)	398	19.1 (51)	648	5.1 (41)
0851	882	52.9 (147)	812	29.6 (129)	295	22.4 (97)	319	20.4 (57)	492	6.9 (69)
0901	1,955	50.0 (129)	1,947	28.2 (118)	650	34.0 (43)	258	22.1 (68)	580	6.0 (55)
0902	74	* *	83	* *	22	* *	6	* *	23	* *
1051	1,549	46.2 (105)	1,470	24.6 (89)	417	36.5 (36)	150	14.7 (30)	437	4.1 (22)
1055	1	* *	1	* *	0	* *	0	* *	0	* *
1101	1,742	37.9 (64)	1,919	23.4 (78)	431	29.0 (73)	509	22.4 (72)	867	6.8 (67)
1102	934	49.6 (125)	908	30.6 (138)	333	29.1 (71)	228	27.2 (80)	410	8.0 (84)
1106	133	49.6 (125)	130	31.5 (142)	49	* *	26	* *	48	* *
1250	1,117	40.9 (75)	1,060	26.3 (103)	308	24.0 (90)	299	15.1 (31)	513	5.3 (47)
1251	736	57.5 (163)	789	38.3 (166)	287	20.6 (99)	243	21.0 (62)	389	7.2 (73)
1252	619	43.9 (87)	584	25.2 (92)	134	22.4 (97)	210	16.2 (38)	317	4.1 (22)
1253	658	57.9 (164)	675	37.0 (162)	208	18.3 (101)	229	21.0 (62)	423	6.1 (56)
1350	2,689	44.9 (93)	2,664	27.1 (108)	952	27.6 (80)	901	19.2 (53)	1,347	4.9 (36)
1351	154	39.6 (68)	161	23.6 (80)	52	* *	65	* *	94	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

(1)

Page: 133
Run Date: 7 / 21 / 10

Clinic	Comparison of Smoking Indicators (2, 3)									
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal	Postpartum			
Clinic	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1352	495	47.3 (110)	460	32.8 (148)	193	23.8 (91)	159	32.1 (86)	221	10.4 (102)
1401	253	45.1 (97)	243	27.2 (109)	69	* *	22	* *	61	* *
1402	178	37.6 (61)	178	24.2 (86)	43	* *	17	* *	57	* *
1403	30	* *	27	* *	8	* *	2	* *	4	* *
1404	23	* *	11	* *	0	* *	3	* *	6	* *
1501	1,169	29.3 (40)	1,174	19.6 (59)	258	26.7 (83)	307	14.3 (27)	483	6.2 (59)
1502	13	* *	6	* *	4	* *	0	* *	0	* *
1503	1,285	37.0 (59)	1,185	20.7 (66)	306	30.7 (63)	359	15.3 (33)	583	5.1 (41)
1504	2,633	28.1 (38)	2,619	14.7 (38)	538	45.5 (7)	710	5.6 (2)	1,191	1.6 (2)
1505	871	30.9 (43)	918	14.8 (39)	205	36.1 (39)	252	7.1 (4)	419	1.7 (3)
1506	658	42.4 (81)	675	22.2 (71)	219	31.5 (57)	187	20.3 (55)	292	5.1 (41)
1507	710	45.2 (98)	711	22.2 (71)	239	38.9 (28)	180	15.6 (36)	306	5.9 (54)
1508	192	1.6 (2)	142	4.9 (6)	1	* *	34	* *	45	* *
1509	665	30.5 (42)	679	20.3 (61)	155	32.9 (50)	179	8.4 (8)	316	2.8 (9)
1510	228	48.2 (115)	215	32.1 (145)	85	* *	35	* *	56	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

(1)

Page: 134
Run Date: 7 / 21 / 10

Clinic	Comparison of Smoking Indicators (2,3)									
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal	Postpartum			
Clinic	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
1511	647	46.1 (104)	670	30.1 (133)	233	29.2 (70)	192	9.9 (14)	319	4.1 (22)
1512	284	44.7 (91)	300	24.3 (87)	92	* *	68	* *	129	7.0 (70)
1701	684	55.3 (157)	689	29.0 (122)	232	25.9 (85)	46	* *	195	6.7 (66)
1712	91	* *	83	* *	38	* *	6	* *	14	* *
1950	726	45.9 (101)	712	27.4 (113)	221	27.6 (80)	176	27.3 (82)	305	7.5 (80)
1952	163	51.5 (139)	173	28.9 (121)	52	* *	24	* *	56	* *
1953	360	50.3 (132)	351	26.8 (106)	110	38.2 (30)	65	* *	124	4.0 (20)
1954	683	44.9 (93)	662	27.5 (114)	198	31.3 (58)	179	9.5 (11)	299	8.4 (89)
1955	159	44.0 (89)	151	23.8 (83)	45	* *	32	* *	59	* *
1957	463	45.4 (99)	487	22.8 (76)	146	37.0 (33)	103	20.4 (57)	201	3.5 (16)
1959	212	53.8 (150)	217	20.3 (61)	79	* *	40	* *	65	* *
1963	35	* *	25	* *	11	* *	0	* *	0	* *
2150	732	55.2 (155)	730	26.0 (101)	227	40.1 (23)	54	* *	205	3.4 (15)
2151	292	59.6 (167)	302	34.4 (156)	108	27.8 (79)	30	* *	80	* *
2251	494	54.5 (151)	493	30.8 (140)	158	31.6 (56)	35	* *	123	8.9 (92)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

(1)

Page: 135
Run Date: 7 / 21 / 10

Clinic	Comparison of Smoking Indicators (2, 3)											
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household						
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal	Postpartum					
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
2252	261	53.3 (149)	251	30.3 (134)	91	* *	22	* *	64	* *		
2551	4,160	36.3 (57)	3,158	20.6 (65)	687	33.8 (46)	513	19.1 (51)	476	6.5 (64)		
2552	8	* *	0	* *	0	* *	0	* *	0	* *		
2554	122	48.4 (118)	116	29.3 (126)	33	* *	13	* *	26	* *		
2555	4,265	44.8 (92)	3,642	27.0 (107)	886	34.3 (42)	617	19.4 (54)	803	7.3 (75)		
3001	974	35.0 (53)	983	21.4 (70)	207	30.9 (61)	117	27.4 (83)	312	7.4 (78)		
3002	249	31.7 (46)	251	20.7 (66)	46	* *	32	* *	78	* *		
3003	6	* *	7	* *	0	* *	1	* *	4	* *		
3004	213	4.7 (6)	195	1.5 (1)	6	* *	24	* *	45	* *		
3007	31	* *	33	* *	1	* *	2	* *	7	* *		
3009	9	* *	11	* *	0	* *	0	* *	3	* *		
3010	1	* *	1	* *	0	* *	0	* *	0	* *		
3201	604	45.0 (95)	626	25.9 (100)	175	28.0 (77)	78	* *	181	6.1 (56)		
3301	4,196	38.2 (66)	4,222	19.8 (60)	1,183	36.3 (37)	1,608	14.3 (27)	2,394	4.5 (31)		
3302	297	34.0 (51)	317	20.5 (63)	69	* *	99	* *	155	3.2 (13)		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

(1)

Page: 136
Run Date: 7 / 21 / 10

Clinic	Comparison of Smoking Indicators (2, 3)											
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household						
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum				
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)		
3303	15	* *	16	* *	8	* *	0	* *	0	* *		
3304	101	4.0 (3)	113	3.5 (4)	3	* *	31	* *	52	* *		
3305	100	34.0 (51)	119	16.8 (48)	29	* *	27	* *	57	* *		
3306	161	46.0 (102)	186	28.0 (117)	54	* *	60	* *	113	6.2 (59)		
3308	54	* *	48	* *	17	* *	14	* *	21	* *		
3309	37	* *	42	* *	14	* *	11	* *	18	* *		
3311	13	* *	10	* *	2	* *	0	* *	0	* *		
3550	80	* *	82	* *	35	* *	11	* *	23	* *		
3551	35	* *	33	* *	15	* *	2	* *	6	* *		
3552	46	* *	52	* *	23	* *	0	* *	11	* *		
3701	987	43.9 (87)	1,005	26.7 (104)	279	25.8 (87)	231	23.8 (74)	429	6.3 (61)		
3702	132	45.5 (100)	114	33.3 (150)	46	* *	0	* *	0	* *		
3703	57	* *	50	* *	19	* *	0	* *	0	* *		
3704	55	* *	49	* *	17	* *	0	* *	0	* *		
3805	3,724	49.5 (124)	3,583	30.4 (136)	1,460	29.1 (71)	1,249	21.1 (64)	1,848	6.3 (61)		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

(1)

Page: 137
Run Date: 7 / 21 / 10

Clinic	Comparison of Smoking Indicators (2,3)									
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal	Postpartum			
	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
3950	2,310	39.7 (69)	2,272	23.0 (77)	660	33.8 (46)	535	15.1 (31)	898	4.8 (35)
3952	45	* *	48	* *	18	* *	2	* *	5	* *
3953	38	* *	35	* *	8	* *	1	* *	5	* *
3954	128	42.2 (80)	125	15.2 (41)	39	* *	15	* *	27	* *
4001	1,210	35.0 (53)	1,120	20.9 (68)	283	27.9 (78)	281	20.3 (55)	480	11.9 (109)
4002	978	40.3 (71)	970	22.7 (75)	268	31.7 (55)	241	27.0 (79)	402	14.2 (115)
4103	105	39.0 (67)	112	20.5 (63)	27	* *	14	* *	46	* *
4106	1,055	26.6 (35)	1,090	13.2 (31)	195	39.0 (27)	182	11.5 (16)	362	5.2 (45)
4107	3,603	35.3 (55)	3,628	16.8 (48)	881	40.0 (24)	1,006	12.8 (23)	1,721	4.4 (29)
4109	1,273	42.7 (84)	1,316	23.5 (79)	400	34.0 (43)	294	22.1 (68)	547	4.9 (36)
4110	1,917	28.7 (39)	1,912	15.8 (44)	367	30.5 (66)	473	12.7 (22)	778	4.1 (22)
4111	1	* *	0	* *	0	* *	0	* *	0	* *
4112	795	1.4 (1)	756	1.9 (2)	9	* *	174	6.3 (3)	280	4.3 (28)
4113	976	13.8 (11)	897	7.6 (11)	100	44.0 (8)	166	2.4 (1)	300	1.7 (3)
4120	2,177	33.8 (50)	2,279	17.1 (51)	540	38.3 (29)	531	12.2 (19)	934	2.0 (5)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

(1)

Page: 138
Run Date: 7 / 21 / 10

Clinic	Comparison of Smoking Indicators (2,3)									
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
Clinic	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4160	2,022	27.5 (37)	2,073	12.9 (30)	385	41.6 (13)	495	7.9 (6)	816	2.1 (6)
4351	712	48.0 (113)	698	31.1 (141)	224	26.8 (82)	71	* *	188	9.6 (98)
4352	192	55.2 (155)	184	37.5 (165)	60	* *	21	* *	51	* *
4353	589	42.4 (81)	572	25.7 (97)	149	36.2 (38)	57	* *	153	6.5 (64)
4354	866	50.2 (131)	850	29.2 (125)	263	31.2 (60)	101	20.8 (60)	250	10.8 (103)
4355	402	56.5 (159)	410	32.4 (147)	150	31.3 (58)	43	* *	100	5.0 (39)
4356	603	48.3 (116)	615	26.2 (102)	173	35.3 (41)	69	* *	163	4.9 (36)
4357	16	* *	17	* *	1	* *	0	* *	2	* *
4358	232	4.3 (4)	207	2.4 (3)	6	* *	34	* *	66	* *
4359	525	40.4 (73)	519	25.8 (98)	108	20.4 (100)	56	* *	165	3.6 (17)
4360	195	49.2 (122)	185	29.7 (130)	52	* *	15	* *	42	* *
4361	282	60.3 (168)	274	41.6 (170)	106	23.6 (93)	29	* *	78	* *
4362	378	52.1 (143)	387	36.7 (161)	138	25.4 (88)	29	* *	109	11.0 (104)
4363	1,035	50.4 (133)	1,027	28.8 (120)	317	31.9 (53)	110	20.9 (61)	270	7.4 (78)
4401	891	46.5 (107)	1,418	29.0 (122)	256	29.3 (69)	144	22.2 (70)	417	7.9 (82)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

(1)

Page: 139
Run Date: 7 / 21 / 10

Clinic	Comparison of Smoking Indicators (2, 3)									
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)		Smoking in Household			
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy				Prenatal		Postpartum	
Clinic	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
4501	1,075	46.6 (108)	1,088	25.4 (95)	324	37.0 (33)	117	15.4 (34)	308	4.2 (27)
4601	1,526	37.7 (63)	1,669	23.6 (80)	389	31.9 (53)	316	11.1 (15)	580	5.2 (45)
4701	1,156	51.0 (135)	1,395	25.8 (98)	413	40.4 (20)	236	17.4 (44)	514	5.8 (52)
5050	1,344	41.9 (79)	450	14.0 (35)	78	* *	248	16.9 (41)	176	4.0 (20)
5051	2,761	38.0 (65)	1,057	16.1 (45)	182	48.4 (4)	465	14.4 (29)	401	5.5 (49)
5052	2,032	32.5 (48)	634	14.5 (36)	88	* *	414	12.1 (18)	266	3.0 (11)
5054	1	* *	0	* *	0	* *	0	* *	0	* *
5101	1,421	36.9 (58)	1,729	19.1 (57)	303	40.3 (22)	370	14.1 (25)	822	4.5 (31)
5202	584	51.4 (138)	640	27.3 (111)	200	33.5 (49)	60	* *	186	3.2 (13)
5203	200	56.5 (159)	212	32.1 (145)	66	* *	13	* *	43	* *
5204	189	46.0 (102)	180	21.1 (69)	51	* *	10	* *	47	* *
5205	31	* *	33	* *	9	* *	0	* *	0	* *
5601	1,213	48.4 (118)	1,266	25.2 (92)	359	34.0 (43)	145	18.6 (48)	380	4.5 (31)
5801	2,394	48.3 (116)	2,357	29.0 (122)	766	32.2 (51)	478	18.8 (49)	890	7.9 (82)
5803	2	* *	22	* *	0	* *	1	* *	18	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

(1)

Page: 140
Run Date: 7 / 21 / 10

Clinic	Comparison of Smoking Indicators (2, 3)											
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household						
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal	Postpartum					
	Number	%	(Rank)	Number	%	(Rank)	Number	%	(Rank)	Number	%	(Rank)
5804	0	*	*	11	*	*	0	*	*	7	*	*
6101	44	*	*	42	*	*	16	*	*	11	*	*
6102	133	42.9	(85)	163	23.9	(84)	43	*	*	44	*	*
6103	4,640	41.1	(76)	4,621	24.5	(88)	1,344	32.1	(52)	1,012	16.1	(37)
6104	61	*	*	71	*	*	21	*	*	24	*	*
6105	39	*	*	43	*	*	16	*	*	13	*	*
6106	1	*	*	5	*	*	0	*	*	1	*	*
6301	4,703	31.3	(44)	5,143	15.2	(41)	953	42.6	(10)	812	11.9	(17)
6312	2,603	20.8	(27)	2,818	8.6	(16)	342	55.8	(2)	417	9.8	(13)
6316	1,302	35.6	(56)	1,483	17.9	(52)	308	41.6	(13)	230	8.7	(9)
6320	257	23.0	(31)	268	15.7	(43)	28	*	*	36	*	*
6325	160	47.5	(111)	151	27.2	(109)	36	*	*	21	*	*
6330	470	29.4	(41)	440	17.0	(50)	76	*	*	88	*	*
6335	622	22.3	(30)	638	13.3	(32)	81	*	*	104	8.7	(9)
6340	200	26.0	(34)	168	17.9	(52)	23	*	*	17	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

(1)

Page: 141
Run Date: 7 / 21 / 10

Clinic	Comparison of Smoking Indicators (2, 3)											
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household						
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal		Postpartum				
	Number	%	(Rank)	Number	%	(Rank)	Number	%	(Rank)	Number	%	(Rank)
6345	52	*	*	52	*	*	13	*	*	7	*	*
6350	276	37.0	(59)	266	22.6	(73)	60	*	*	45	*	*
7304	238	40.3	(71)	244	24.6	(89)	48	*	*	24	*	*
7306	2,639	32.4	(47)	2,586	17.9	(52)	470	41.5	(15)	371	16.7	(39)
7308	210	40.5	(74)	211	23.7	(82)	51	*	*	32	*	*
7310	24	*	*	33	*	*	10	*	*	6	*	*
7401	2,315	51.2	(136)	1,904	30.6	(138)	659	33.7	(48)	358	21.2	(65)
7402	277	58.5	(166)	218	33.9	(155)	81	*	*	22	*	*
7403	109	45.0	(95)	86	*	*	28	*	*	5	*	*
7404	108	42.6	(83)	87	*	*	24	*	*	13	*	*
7601	793	47.9	(112)	795	29.7	(130)	255	29.8	(68)	100	17.0	(42)
7850	729	52.1	(143)	1,045	33.7	(152)	248	23.0	(96)	195	24.1	(75)
7851	179	52.0	(141)	194	37.1	(164)	61	*	*	39	*	*
7852	100	40.0	(70)	119	30.3	(134)	30	*	*	16	*	*
8150	66	*	*	70	*	*	10	*	*	9	*	*
										20	*	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

(1)

Page: 142
Run Date: 7 / 21 / 10

Clinic	Comparison of Smoking Indicators (2, 3)									
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal	Postpartum			
Clinic	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8151	2,884	31.4 (45)	2,879	16.7 (47)	520	35.8 (40)	624	17.8 (45)	1,129	8.1 (86)
8152	560	46.8 (109)	541	22.6 (73)	200	40.5 (19)	107	37.4 (88)	188	13.8 (113)
8153	38	* *	37	* *	12	* *	4	* *	9	* *
8154	306	10.5 (8)	304	5.3 (7)	12	* *	39	* *	94	* *
8155	93	* *	91	* *	8	* *	8	* *	20	* *
8156	44	* *	41	* *	8	* *	5	* *	9	* *
8157	24	* *	22	* *	4	* *	11	* *	10	* *
8201	2,089	37.6 (61)	1,450	18.1 (55)	244	42.6 (10)	491	12.2 (19)	616	2.8 (9)
8202	2,419	14.1 (15)	2,295	10.8 (26)	123	40.7 (18)	542	7.9 (6)	880	3.8 (19)
8203	201	46.3 (106)	159	42.8 (171)	20	* *	74	* *	54	* *
8250	252	27.4 (36)	226	13.3 (32)	29	* *	247	22.7 (73)	219	4.1 (22)
8255	2,623	41.2 (77)	1,580	16.6 (46)	314	43.3 (9)	520	17.9 (46)	331	5.1 (41)
8257	344	14.2 (16)	308	7.1 (10)	12	* *	114	12.3 (21)	132	8.3 (88)
8258	1,360	41.7 (78)	859	34.5 (157)	151	40.4 (20)	255	14.1 (25)	329	3.0 (11)
8260	413	16.7 (21)	453	8.4 (13)	41	* *	86	* *	163	6.1 (56)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

(1)

Page: 143
Run Date: 7 / 21 / 10

Clinic	Comparison of Smoking Indicators (2, 3)									
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal	Postpartum			
Clinic	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
8901	846	19.3 (26)	991	10.9 (27)	92	* *	148	25.7 (76)	298	11.4 (107)
8902	481	15.0 (19)	608	9.0 (19)	41	* *	95	* *	213	11.7 (108)
8903	1,048	16.8 (22)	1,195	8.6 (16)	106	41.5 (15)	202	21.3 (66)	398	10.3 (101)
8904	771	13.7 (10)	891	8.4 (13)	65	* *	139	25.9 (77)	289	10.0 (99)
8905	1,853	14.6 (17)	2,399	13.5 (34)	149	30.9 (61)	458	9.6 (12)	914	7.2 (73)
9004	306	16.3 (20)	273	9.2 (20)	29	* *	3	* *	3	* *
9010	34	* *	29	* *	2	* *	0	* *	0	* *
9012	10	* *	15	* *	0	* *	10	* *	12	* *
9013	12	* *	34	* *	1	* *	12	* *	32	* *
9015	9	* *	11	* *	0	* *	9	* *	10	* *
9016	21	* *	36	* *	1	* *	21	* *	31	* *
9017	8	* *	10	* *	0	* *	8	* *	10	* *
9018	19	* *	34	* *	2	* *	19	* *	31	* *
9019	10	* *	13	* *	0	* *	10	* *	12	* *
9021	14	* *	22	* *	2	* *	14	* *	20	* *

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

(1)

Page: 144
Run Date: 7 / 21 / 10

Clinic	Comparison of Smoking Indicators (2, 3)									
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household				
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal	Postpartum			
Clinic	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
9023	59	* *	52	* *	5	* *	0	* *	0	* *
9040	243	23.0 (31)	220	11.4 (28)	36	* *	3	* *	6	* *
9041	718	12.3 (9)	675	8.9 (18)	35	* *	41	* *	148	9.5 (97)
9045	1,247	8.3 (7)	1,243	9.8 (23)	59	* *	233	15.5 (35)	441	7.3 (75)
9050	1,947	13.8 (11)	2,061	6.3 (9)	166	48.2 (5)	393	16.8 (40)	727	8.1 (86)
9051	2,377	13.9 (13)	2,483	8.3 (12)	155	37.4 (31)	392	7.1 (4)	844	2.7 (8)
9053	2,289	24.9 (33)	2,579	14.6 (37)	286	40.9 (17)	505	21.4 (67)	995	13.4 (112)
9054	78	* *	79	* *	5	* *	0	* *	0	* *
9055	77	* *	105	12.4 (29)	6	* *	2	* *	21	* *
9058	541	4.3 (4)	527	3.8 (5)	9	* *	145	17.2 (43)	236	6.4 (63)
9066	2,417	14.0 (14)	2,604	8.5 (15)	200	39.5 (25)	441	20.6 (59)	854	10.2 (100)
9068	919	32.5 (48)	1,294	18.9 (56)	160	30.6 (64)	254	28.0 (84)	532	14.5 (116)
9069	2,166	18.9 (24)	2,358	9.2 (20)	262	54.2 (3)	552	25.9 (77)	1,034	8.9 (92)
9070	1,713	21.2 (28)	1,935	10.1 (24)	194	37.1 (32)	353	13.9 (24)	766	8.5 (90)
9072	1,384	18.5 (23)	1,716	15.1 (40)	141	41.8 (12)	371	18.1 (47)	683	11.1 (105)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 7A
2007 - 2009 Pregnancy Nutrition Surveillance
Michigan
Comparison of Smoking Indicators by Clinic
Combined 3 Years

(1)

Page: 145
Run Date: 7 / 21 / 10

Clinic	Comparison of Smoking Indicators (2, 3)												
	Smoking				Quit Smoking By First Prenatal Visit and Stayed Off Cigarettes (4)	Smoking in Household							
	3 Months Prior to Pregnancy		Last 3 Months of Pregnancy			Prenatal	Postpartum						
Clinic	Number	%	(Rank)	Number	%	(Rank)	Number	%	(Rank)	Number	%	(Rank)	
9074	1,601	22.1	(29)	1,668	10.4	(25)	242	56.2	(1)	345	27.2	(80)	
9079	533	14.6	(17)	442	5.9	(8)	39	*	*	0	*	*	
9080	2,215	19.2	(25)	2,246	9.6	(22)	280	47.1	(6)	418	22.2	(70)	
Unknown Clinics	8	*	*	240	93.3	(172)	0	*	*	8	*	*	
Michigan	166,382	36.2		163,164	20.5		36,412	34.1		34,277	16.9		
												60,783	6.4

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) Among women who smoked 3 months prior to pregnancy.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Page: 146
Run Date: 7 / 21 / 10

Clinic	Comparison of Infant Health Indicators (2, 3)							
	Birthweight		Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)	High (5)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0101	132	7.6 (98) 10.6 (149)	143	8.4 (167)	122	4.9 (67)	138	60.1 (86)
0102	406	7.1 (83) 10.3 (145)	436	4.8 (58)	382	3.9 (23)	418	51.9 (129)
0103	414	4.6 (12) 11.4 (161)	450	2.4 (6)	404	3.7 (18)	426	52.6 (126)
0104	130	8.5 (128) 7.7 (80)	135	4.4 (39)	123	7.3 (141)	132	50.8 (132)
0250	148	6.1 (43) 10.8 (153)	152	5.9 (116)	138	4.3 (38)	146	65.1 (53)
0251	116	4.3 (7) 9.5 (127)	144	4.2 (30)	110	4.5 (47)	137	51.1 (130)
0252	129	5.4 (17) 13.2 (170)	128	5.5 (91)	120	2.5 (6)	121	69.4 (27)
0253	153	9.8 (148) 11.8 (166)	165	4.2 (30)	144	8.3 (150)	150	54.0 (119)
0401	550	8.0 (112) 8.2 (96)	577	4.3 (33)	523	4.8 (62)	554	50.0 (134)
0402	464	3.9 (5) 10.8 (153)	486	3.7 (23)	447	2.7 (7)	468	67.7 (36)
0403	122	9.0 (134) 7.4 (70)	127	3.1 (14)	118	7.6 (145)	122	61.5 (80)
0404	139	5.8 (33) 7.9 (87)	147	4.8 (58)	133	4.5 (47)	140	62.9 (68)
0501	528	5.5 (22) 9.7 (132)	570	3.9 (26)	507	4.3 (38)	526	69.0 (28)
0502	4	* * * *	4	*	4	*	4	*

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Page: 147
Run Date: 7 / 21 / 10

Clinic	Comparison of Infant Health Indicators (2, 3)							
	Birthweight		Preterm (6)		Full Term LBW (7)		Ever Breastfed	
	Low (4)	High (5)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)
0503	166	10.8 (153) 10.8 (153)	178	6.7 (138)	154	8.4 (153)	166	79.5 (9)
0504	569	6.7 (69) 9.0 (120)	608	4.4 (39)	542	4.4 (42)	575	76.2 (13)
0505	349	5.7 (30) 8.3 (102)	384	2.6 (9)	339	5.0 (75)	358	72.9 (19)
0506	185	9.7 (147) 7.0 (56)	193	4.7 (51)	176	6.8 (136)	185	70.3 (24)
0650	992	7.2 (87) 7.8 (84)	1,134	6.8 (144)	910	5.7 (110)	1,061	67.1 (39)
0651	617	9.4 (141) 5.2 (29)	663	5.6 (99)	579	8.5 (154)	617	54.6 (112)
0652	325	7.4 (93) 7.4 (70)	348	4.3 (33)	312	5.1 (80)	320	56.9 (101)
0653	419	6.0 (40) 7.4 (70)	444	5.4 (87)	395	4.1 (30)	429	64.1 (59)
0654	356	7.6 (98) 11.5 (163)	386	4.1 (29)	336	5.4 (99)	355	67.0 (40)
0655	410	6.8 (71) 6.6 (49)	444	4.7 (51)	388	5.2 (90)	415	66.3 (44)
0657	68	* * * *	67	* *	66	* *	67	* *
0701	404	3.7 (4) 9.7 (132)	435	5.1 (73)	384	2.9 (9)	398	76.1 (14)
0702	97	* * * *	100	7.0 (152)	91	* *	90	* *
0703	255	7.1 (83) 9.8 (134)	258	5.8 (108)	239	5.4 (99)	243	61.7 (77)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Page: 148
Run Date: 7 / 21 / 10

Clinic	Comparison of Infant Health Indicators (2, 3)									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)
0704	76	* * * *	79	* *	73	* *	74	* *		
0705	171	2.3 (1) 9.9 (137)	183	2.7 (10)	165	1.8 (2)	172	65.1 (53)		
0850	1,166	6.1 (43) 9.1 (123)	1,218	5.7 (106)	1,094	4.0 (25)	1,172	67.2 (38)		
0851	926	5.8 (33) 9.0 (120)	952	5.8 (108)	873	4.0 (25)	928	56.3 (104)		
0901	1,886	7.3 (88) 10.7 (152)	1,946	5.1 (73)	1,783	5.3 (96)	1,889	54.4 (114)		
0902	81	* * * *	83	* *	78	* *	79	* *		
1051	1,429	6.2 (47) 10.2 (143)	1,513	5.8 (108)	1,345	4.4 (42)	1,423	81.2 (7)		
1055	1	* * * *	2	* *	1	* *	1	* *		
1101	1,858	8.2 (120) 6.5 (45)	1,940	5.8 (108)	1,746	5.8 (112)	1,837	45.8 (144)		
1102	889	7.4 (93) 9.8 (134)	913	5.6 (99)	835	5.1 (80)	881	54.3 (115)		
1106	130	5.4 (17) 7.7 (80)	134	6.7 (138)	121	4.1 (30)	131	64.1 (59)		
1250	1,033	5.5 (22) 8.3 (102)	1,089	4.9 (63)	974	3.7 (18)	1,055	61.0 (83)		
1251	874	6.5 (58) 6.2 (40)	909	4.3 (33)	832	5.2 (90)	877	62.5 (72)		
1252	594	5.6 (26) 6.2 (40)	702	5.3 (83)	559	4.3 (38)	673	64.2 (58)		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Page: 149
Run Date: 7 / 21 / 10

Clinic	Comparison of Infant Health Indicators (2, 3)									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)	Number (%) (Rank)							
1253	742	6.1 (43) 7.4 (70)	802	6.9 (147)	686	3.2 (14)	782	56.6 (102)		
1350	2,613	7.4 (93) 8.2 (96)	2,680	4.9 (63)	2,478	5.2 (90)	2,595	51.0 (131)		
1351	156	5.1 (14) 7.1 (58)	159	6.9 (147)	145	4.1 (30)	155	67.7 (36)		
1352	437	6.9 (76) 7.8 (84)	449	4.7 (51)	414	4.8 (62)	438	49.3 (137)		
1401	225	9.3 (139) 10.2 (143)	240	4.6 (47)	213	7.0 (139)	226	68.6 (33)		
1402	178	5.1 (14) 14.0 (171)	179	3.4 (19)	172	4.1 (30)	177	83.6 (6)		
1403	25	* * * *	27	* *	25	* *	25	* *		
1404	12	* * * *	12	* *	10	* *	12	* *		
1501	1,066	6.5 (58) 8.2 (96)	1,228	4.8 (58)	1,006	5.1 (80)	1,138	62.2 (75)		
1502	10	* * * *	11	* *	10	* *	10	* *		
1503	1,290	6.4 (53) 8.8 (118)	1,425	5.7 (106)	1,212	5.2 (90)	1,302	63.7 (62)		
1504	2,383	6.6 (60) 7.2 (60)	2,679	5.4 (87)	2,211	4.8 (62)	2,557	62.8 (70)		
1505	853	6.3 (48) 10.1 (141)	908	5.5 (91)	804	4.2 (37)	880	68.3 (35)		
1506	637	6.6 (60) 10.0 (140)	686	5.5 (91)	601	4.8 (62)	649	63.2 (64)		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Page: 150
Run Date: 7 / 21 / 10

Clinic	Comparison of Infant Health Indicators (2, 3)									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)	Number (%) (Rank)							
1507	668	6.3 (48) 11.1 (159)	718	6.4 (131)	626	4.6 (55)	687	65.9 (45)		
1508	113	8.0 (112) 8.0 (88)	183	2.7 (10)	109	8.3 (150)	164	64.6 (57)		
1509	649	5.7 (30) 8.2 (96)	720	6.3 (128)	605	3.8 (21)	690	58.3 (94)		
1510	208	8.2 (120) 10.6 (149)	222	3.6 (21)	200	6.5 (132)	214	50.0 (134)		
1511	645	6.7 (69) 11.5 (163)	660	5.9 (116)	607	4.0 (25)	643	63.1 (65)		
1512	289	6.6 (60) 11.4 (161)	302	5.6 (99)	275	4.4 (42)	287	58.2 (95)		
1701	658	5.9 (35) 13.1 (169)	693	3.9 (26)	627	4.3 (38)	649	64.7 (56)		
1712	73	* * * *	82	* *	69	* *	77	* *		
1950	677	6.4 (53) 7.5 (77)	730	6.2 (122)	630	4.1 (30)	689	60.4 (85)		
1952	152	6.6 (60) 10.5 (148)	178	6.2 (122)	141	2.8 (8)	164	68.9 (30)		
1953	327	8.0 (112) 8.6 (114)	349	5.2 (76)	307	5.2 (90)	331	59.5 (90)		
1954	620	7.9 (108) 7.3 (66)	670	7.5 (159)	569	5.4 (99)	632	62.7 (71)		
1955	135	4.4 (8) 11.1 (159)	152	5.3 (83)	129	3.1 (13)	139	63.3 (63)		
1957	461	5.4 (17) 7.4 (70)	477	6.5 (133)	429	3.3 (15)	454	65.2 (51)		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Page: 151
Run Date: 7 / 21 / 10

Clinic	Comparison of Infant Health Indicators (2, 3)									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)								
	Number	% (Rank)	% (Rank)	Number	% (Rank)	Number	% (Rank)	Number	% (Rank)	
1959	204	6.9 (76)	10.8 (153)	210	5.2 (76)	191	5.8 (112)	197	73.1 (18)	
1963	23	*	*	23	*	23	*	22	*	
2150	716	5.7 (30)	10.8 (153)	741	4.9 (63)	681	4.6 (55)	698	59.7 (88)	
2151	304	7.6 (98)	7.2 (60)	318	5.0 (69)	289	5.9 (119)	293	47.8 (140)	
2251	471	6.6 (60)	10.4 (147)	489	4.5 (45)	449	4.5 (47)	471	59.7 (88)	
2252	235	6.0 (40)	8.1 (90)	247	6.9 (147)	218	4.1 (30)	236	60.6 (84)	
2551	3,152	11.2 (156)	4.3 (12)	3,261	4.3 (33)	3,001	9.4 (161)	3,201	39.6 (155)	
2552	0	*	*	0	*	0	*	0	*	
2554	111	5.4 (17)	8.1 (90)	116	3.4 (19)	108	4.6 (55)	110	30.9 (163)	
2555	3,683	7.8 (105)	7.2 (60)	3,904	4.4 (39)	3,492	6.2 (125)	3,801	53.4 (122)	
3001	994	11.8 (162)	3.7 (4)	1,032	7.2 (157)	924	9.5 (163)	937	41.3 (153)	
3002	235	11.1 (155)	6.8 (53)	247	8.1 (165)	218	8.3 (150)	220	33.2 (162)	
3003	9	*	*	9	*	9	*	9	*	
3004	183	5.5 (22)	4.4 (15)	187	5.3 (83)	173	4.0 (25)	178	65.2 (51)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Page: 152
Run Date: 7 / 21 / 10

Clinic	Comparison of Infant Health Indicators (2, 3)									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)	Number % (Rank)	% (Rank)	Number % (Rank)	% (Rank)	Number % (Rank)	% (Rank)		
3007	31	* * * *	35	* *	30	* *	30	* *		
3009	13	* * * *	14	* *	13	* *	13	* *		
3010	1	* * * *	2	* *	1	* *	1	* *		
3201	574	5.6 (26) 11.0 (158)	608	5.4 (87)	540	3.9 (23)	579	54.2 (118)		
3301	4,121	7.3 (88) 7.3 (66)	4,492	5.8 (108)	3,860	4.9 (67)	4,283	63.1 (65)		
3302	300	8.7 (131) 5.7 (35)	332	6.9 (147)	282	7.4 (143)	306	52.0 (128)		
3303	15	* * * *	17	* *	15	* *	15	* *		
3304	109	6.4 (53) 4.6 (17)	117	4.3 (33)	103	2.9 (9)	116	99.1 (1)		
3305	111	4.5 (9) 11.7 (165)	119	2.5 (8)	109	3.7 (18)	113	77.9 (10)		
3306	182	8.8 (133) 9.9 (137)	195	7.2 (157)	167	6.0 (123)	182	66.5 (42)		
3308	53	* * * *	54	* *	50	* *	50	* *		
3309	39	* * * *	41	* *	37	* *	40	* *		
3311	12	* * * *	12	* *	12	* *	12	* *		
3550	76	* * * *	78	* *	74	* *	75	* *		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Page: 153
Run Date: 7 / 21 / 10

Clinic	Comparison of Infant Health Indicators (2, 3)									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)
3551	34	* * * *	34	* *	34	* *	34	* *		
3552	52	* * * *	52	* *	50	* *	51	* *		
3701	973	6.9 (76) 7.3 (66)	1,002	7.1 (153)	901	5.0 (75)	953	55.5 (110)		
3702	115	3.5 (2) 8.7 (116)	117	0.0 (1)	115	3.5 (16)	113	48.7 (139)		
3703	51	* * * *	51	* *	51	* *	50	* *		
3704	50	* * * *	50	* *	50	* *	48	* *		
3805	3,546	7.1 (83) 8.5 (110)	3,607	5.2 (76)	3,342	5.1 (80)	3,469	54.5 (113)		
3950	2,228	8.1 (117) 8.3 (102)	2,272	5.3 (83)	2,101	5.6 (107)	2,209	58.1 (96)		
3952	47	* * * *	49	* *	47	* *	48	* *		
3953	34	* * * *	35	* *	34	* *	34	* *		
3954	125	6.4 (53) 7.2 (60)	123	2.4 (6)	120	5.0 (75)	123	55.3 (111)		
4001	1,016	10.5 (151) 5.2 (29)	1,087	8.9 (172)	918	6.2 (125)	1,034	55.8 (109)		
4002	924	8.4 (126) 7.7 (80)	969	7.6 (161)	852	5.8 (112)	924	61.7 (77)		
4103	109	18.3 (171) 4.6 (17)	109	10.1 (173)	97	* *	109	45.0 (147)		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Page: 154
Run Date: 7 / 21 / 10

Clinic	Comparison of Infant Health Indicators (2, 3)									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)
4106	1,064	8.7 (131)	4.8 (22)	1,088	4.6 (47)	1,010	6.9 (138)	1,034	54.0 (119)	
4107	3,544	7.9 (108)	8.4 (107)	3,649	6.2 (122)	3,315	4.9 (67)	3,493	62.3 (73)	
4109	1,274	8.2 (120)	9.8 (134)	1,300	5.8 (108)	1,199	5.8 (112)	1,237	62.1 (76)	
4110	1,822	6.0 (40)	7.3 (66)	1,873	5.0 (69)	1,727	4.4 (42)	1,817	58.9 (91)	
4111	0	*	*	*	*	0	*	*	0	*
4112	731	4.1 (6)	10.3 (145)	742	4.3 (33)	698	3.0 (11)	734	75.5 (15)	
4113	877	5.9 (35)	8.4 (107)	886	4.9 (63)	834	3.8 (21)	832	71.0 (21)	
4120	2,204	6.9 (76)	9.5 (127)	2,258	4.4 (39)	2,103	4.9 (67)	2,160	61.7 (77)	
4160	2,026	7.3 (88)	9.0 (120)	2,086	4.7 (51)	1,924	5.6 (107)	2,018	61.1 (82)	
4351	669	6.6 (60)	11.8 (166)	695	3.3 (16)	642	5.5 (102)	666	65.3 (49)	
4352	186	8.1 (117)	6.5 (45)	198	3.0 (13)	180	6.7 (134)	189	56.1 (108)	
4353	540	6.9 (76)	6.7 (51)	563	5.5 (91)	507	5.5 (102)	537	58.7 (92)	
4354	805	6.3 (48)	9.4 (124)	834	4.9 (63)	757	4.5 (47)	816	62.9 (68)	
4355	388	9.5 (143)	9.5 (127)	409	6.1 (121)	364	7.7 (146)	379	65.7 (47)	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Page: 155
Run Date: 7 / 21 / 10

Clinic	Comparison of Infant Health Indicators (2, 3)									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)
4356	593	6.4 (53) 9.6 (130)	608	5.8 (108)	559	4.5 (47)	589	56.2 (105)		
4357	15	* * * *	18	* *	15	* *	18	* *		
4358	192	4.7 (13) 9.9 (137)	219	5.9 (116)	176	2.3 (4)	217	74.2 (16)		
4359	534	5.6 (26) 8.1 (90)	571	4.7 (51)	510	4.1 (30)	524	63.9 (61)		
4360	179	4.5 (9) 10.1 (141)	182	4.9 (63)	168	2.4 (5)	174	69.0 (28)		
4361	261	7.3 (88) 6.5 (45)	269	5.2 (76)	246	4.5 (47)	251	64.9 (55)		
4362	368	7.6 (98) 8.7 (116)	375	5.1 (73)	348	6.0 (123)	363	77.4 (11)		
4363	978	7.1 (83) 9.6 (130)	1,012	4.5 (45)	932	4.9 (67)	964	70.5 (22)		
4401	1,355	7.4 (93) 8.5 (110)	1,409	4.7 (51)	1,282	5.5 (102)	1,345	57.8 (100)		
4501	1,010	6.6 (60) 8.6 (114)	1,059	4.6 (47)	955	4.9 (67)	1,031	61.3 (81)		
4601	1,614	7.3 (88) 7.2 (60)	1,714	4.7 (51)	1,509	5.1 (80)	1,679	57.9 (99)		
4701	1,353	6.6 (60) 10.6 (149)	1,395	4.2 (30)	1,294	4.7 (61)	1,342	66.5 (42)		
5050	479	7.9 (108) 5.4 (31)	492	6.5 (133)	448	5.8 (112)	481	96.3 (2)		
5051	1,134	7.4 (93) 8.5 (110)	1,180	5.2 (76)	1,069	5.3 (96)	1,150	91.9 (3)		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Page: 156
Run Date: 7 / 21 / 10

Clinic	Comparison of Infant Health Indicators (2, 3)									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)
5052	674	9.2 (137) 6.7 (51)	703	6.7 (138)	629	6.4 (130)	678	91.6 (4)		
5054	0	* * * *	0	*	0	*	0	*	*	*
5101	1,696	8.0 (112) 6.4 (44)	1,772	7.1 (153)	1,566	4.9 (67)	1,710	60.0 (87)		
5202	621	6.3 (48) 8.9 (119)	633	6.6 (136)	578	4.5 (47)	616	67.0 (40)		
5203	204	5.4 (17) 11.8 (166)	211	3.8 (25)	196	5.1 (80)	203	63.1 (65)		
5204	180	6.1 (43) 8.3 (102)	182	3.3 (16)	174	4.6 (55)	179	70.4 (23)		
5205	33	* * * *	33	*	33	*	31	*	*	*
5601	1,245	8.0 (112) 8.4 (107)	1,274	5.5 (91)	1,173	5.9 (119)	1,225	68.9 (30)		
5801	2,272	6.9 (76) 7.5 (77)	2,353	6.2 (122)	2,125	4.6 (55)	2,272	53.3 (123)		
5803	23	* * * *	25	*	23	*	24	*	*	*
5804	7	* * * *	9	*	5	*	8	*	*	*
6101	41	* * * *	42	*	40	*	40	*	*	*
6102	154	4.5 (9) 6.5 (45)	154	5.2 (76)	144	1.4 (1)	152	68.4 (34)		
6103	4,442	8.2 (120) 7.1 (58)	4,562	6.2 (122)	4,150	5.8 (112)	4,271	46.6 (142)		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Page: 157
Run Date: 7 / 21 / 10

Clinic	Comparison of Infant Health Indicators (2, 3)									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)	Number (Rank)			Number (Rank)				
6104	66	* * * *	70	*	*	60	*	*	66	* *
6105	42	* * * *	42	*	*	39	*	*	39	* *
6106	5	* * * *	5	*	*	5	*	*	5	* *
6301	5,018	8.2 (120) 6.8 (53)	5,168	5.6 (99)		4,707	5.9 (119)		5,039	46.4 (143)
6312	2,806	9.0 (134) 6.9 (55)	2,868	5.6 (99)		2,642	6.3 (128)		2,808	54.0 (119)
6316	1,446	6.6 (60) 7.8 (84)	1,478	4.4 (39)		1,380	4.6 (55)		1,448	53.2 (124)
6320	264	8.3 (125) 7.6 (79)	283	6.4 (131)		247	5.7 (110)		273	53.1 (125)
6325	148	9.5 (143) 4.7 (20)	154	5.8 (108)		138	5.1 (80)		149	58.4 (93)
6330	438	9.6 (146) 5.7 (35)	458	8.5 (170)		396	5.8 (112)		443	42.0 (151)
6335	601	7.7 (103) 8.2 (96)	643	5.0 (69)		567	6.3 (128)		629	45.5 (145)
6340	166	3.6 (3) 6.6 (49)	180	2.8 (12)		158	1.9 (3)		173	52.6 (126)
6345	52	* * * *	52	*	*	49	*	*	52	* *
6350	256	8.6 (130) 5.1 (28)	272	4.8 (58)		243	7.8 (147)		263	40.7 (154)
7304	244	7.8 (105) 7.0 (56)	251	4.4 (39)		230	5.2 (90)		242	50.4 (133)

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Page: 158
Run Date: 7 / 21 / 10

Clinic	Comparison of Infant Health Indicators (2, 3)										
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed			
	Low (4)	High (5)	Number (Rank)			Number (Rank)					
7306	2,788	9.1 (136)	7.2 (60)	2,855	5.5 (91)	2,627	7.0 (139)	2,681	45.5 (145)		
7308	212	5.2 (16)	9.4 (124)	219	7.8 (162)	195	3.6 (17)	203	58.1 (96)		
7310	32	*	*	*	*	32	*	*	32	*	*
7401	1,874	8.4 (126)	6.3 (42)	1,920	7.1 (153)	1,735	5.0 (75)	1,871	49.7 (136)		
7402	222	6.8 (71)	5.9 (37)	224	3.6 (21)	214	5.1 (80)	223	54.3 (115)		
7403	86	*	*	*	*	80	*	*	87	*	*
7404	93	*	*	*	*	89	*	*	94	*	*
7601	772	6.9 (76)	4.5 (16)	788	5.5 (91)	726	5.6 (107)	756	56.2 (105)		
7850	1,019	6.8 (71)	7.4 (70)	1,037	4.8 (58)	970	4.4 (42)	987	56.6 (102)		
7851	186	8.1 (117)	8.1 (90)	194	4.6 (47)	177	6.2 (125)	185	65.9 (45)		
7852	110	5.5 (22)	5.5 (33)	119	8.4 (167)	100	3.0 (11)	111	70.3 (24)		
8150	62	*	*	*	*	71	*	*	65	*	*
8151	2,565	7.8 (105)	8.3 (102)	2,892	5.9 (116)	2,393	5.5 (102)	2,717	65.3 (49)		
8152	525	9.3 (139)	5.9 (37)	540	5.2 (76)	496	6.7 (134)	516	54.3 (115)		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Page: 159
Run Date: 7 / 21 / 10

Clinic	Comparison of Infant Health Indicators (2, 3)									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)	Number (%) (Rank)							
8153	32 * * * *		35 * *		31 * *		34 * *			
8154	306 5.6 (26) 8.2 (96)		318 3.1 (14)		295 5.1 (80)		310 88.1 (5)			
8155	84 * * * *		91 * *		83 * *		84 * *			
8156	33 * * * *		40 * *		33 * *		38 * *			
8157	18 * * * *		22 * *		18 * *		19 * *			
8201	1,314 7.6 (98) 8.5 (110)		1,506 5.6 (99)		1,232 5.0 (75)		1,310 72.8 (20)			
8202	2,089 5.9 (35) 7.7 (80)		2,504 5.5 (91)		1,958 4.0 (25)		2,322 80.7 (8)			
8203	139 7.9 (108) 9.4 (124)		181 3.3 (16)		132 6.8 (136)		137 76.6 (12)			
8250	223 9.4 (141) 5.4 (31)		225 8.4 (167)		200 5.5 (102)		224 58.0 (98)			
8255	1,466 6.3 (48) 8.0 (88)		1,648 3.7 (23)		1,403 4.9 (67)		1,462 68.8 (32)			
8257	293 6.8 (71) 4.1 (9)		371 5.4 (87)		273 4.8 (62)		344 73.3 (17)			
8258	720 6.8 (71) 8.1 (90)		922 4.0 (28)		681 5.3 (96)		727 70.0 (26)			
8260	440 9.8 (148) 4.8 (22)		448 6.5 (133)		411 8.5 (154)		443 65.5 (48)			
8901	965 12.3 (164) 4.2 (11)		1,010 6.7 (138)		892 9.4 (161)		941 28.3 (165)			

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Page: 160
Run Date: 7 / 21 / 10

Clinic	Comparison of Infant Health Indicators (2, 3)									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)	Number (Rank)			Number (Rank)				
8902	601	11.0 (154)	4.8 (22)	627	7.8 (162)	551	8.5 (154)	598	34.8 (159)	
8903	1,169	12.7 (166)	3.7 (4)	1,210	6.9 (147)	1,075	9.8 (165)	1,170	26.8 (168)	
8904	877	12.8 (167)	5.0 (26)	921	7.1 (153)	811	9.1 (159)	884	49.2 (138)	
8905	2,383	8.5 (128)	7.4 (70)	2,501	6.2 (122)	2,215	6.6 (133)	2,374	36.7 (156)	
9004	293	11.6 (159)	3.4 (2)	297	0.3 (2)	292	11.6 (168)	286	26.6 (169)	
9010	31	*	*	32	*	*	*	*	*	
9012	12	*	*	14	*	*	*	*	*	
9013	32	*	*	33	*	*	26	*	*	
9015	10	*	*	11	*	*	7	*	*	
9016	33	*	*	34	*	*	24	*	*	
9017	10	*	*	10	*	*	10	*	*	
9018	33	*	*	34	*	*	27	*	*	
9019	13	*	*	13	*	*	12	*	*	
9021	20	*	*	21	*	*	17	*	*	

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Page: 161
Run Date: 7 / 21 / 10

Clinic	Comparison of Infant Health Indicators (2, 3)									
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed		
	Low (4)	High (5)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)
9023	67 * * * *		67	* *	67	* *	64	* *		
9040	239 5.9 (35) 3.3 (1)		245	0.4 (3)	238	5.9 (119)	240	23.3 (170)		
9041	685 11.5 (157) 4.1 (9)		783	5.0 (69)	641	10.3 (166)	732	28.0 (166)		
9045	1,261 9.2 (137) 6.3 (42)		1,322	8.0 (164)	1,150	6.4 (130)	1,273	42.0 (151)		
9050	1,957 9.5 (143) 6.1 (39)		2,055	6.8 (144)	1,809	7.5 (144)	1,995	42.3 (150)		
9051	2,455 5.9 (35) 8.1 (90)		2,703	6.0 (120)	2,284	4.5 (47)	2,580	56.2 (105)		
9053	2,765 11.6 (159) 4.3 (12)		2,947	6.3 (128)	2,581	8.9 (157)	2,722	35.1 (158)		
9054	78 * * * *		79	* *	78	* *	79	* *		
9055	105 14.3 (170) 4.8 (22)		108	5.6 (99)	99	* *	105	22.9 (171)		
9058	504 7.7 (103) 5.0 (26)		515	6.6 (136)	467	5.1 (80)	514	62.3 (73)		
9066	2,445 12.0 (163) 3.9 (7)		2,595	6.7 (138)	2,266	9.3 (160)	2,497	27.4 (167)		
9068	1,467 11.7 (161) 4.6 (17)		1,521	7.5 (159)	1,337	8.2 (148)	1,421	34.3 (161)		
9069	2,315 11.5 (157) 4.7 (20)		2,429	6.7 (138)	2,142	9.0 (158)	2,317	43.5 (149)		
9070	1,980 10.5 (151) 3.7 (4)		2,151	6.3 (128)	1,837	8.2 (148)	2,085	34.6 (160)		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.



Table 8A
2007 - 2009 Pregnancy Nutrition Surveillance (1)
Michigan
Comparison of Infant Health Indicators by Clinic
Combined 3 Years

Page: 162
Run Date: 7 / 21 / 10

Clinic	Comparison of Infant Health Indicators (2, 3)											
	Birthweight			Preterm (6)		Full Term LBW (7)		Ever Breastfed				
	Low (4)	High (5)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)	Number (Rank)	% (Rank)		
9072	1,717	13.7 (168)	4.0 (8)		1,840	8.8 (171)	1,555	9.6 (164)	1,765	30.2 (164)		
9074	1,660	13.8 (169)	3.5 (3)		1,696	8.3 (166)	1,512	10.4 (167)	1,607	44.9 (148)		
9079	415	12.3 (164)	4.3 (12)		449	0.4 (3)	413	12.3 (169)	425	36.0 (157)		
9080	2,216	10.0 (150)	5.6 (34)		2,295	6.8 (144)	2,062	7.3 (141)	2,232	47.8 (140)		
Unknown Clinics	18	*	*	*	265	1.5 (5)	18	*	*	87	*	*
Michigan	159,197	8.0	7.4		168,223	5.6	149,422	5.9	160,354	55.3		

(1) Reporting period is January 1 through December 31.

(2) Excludes records with unknown data or errors.

(3) Rank compares one clinic's rate to other clinics. Rank 1 = best rate.

(4) <2500 g.

(5) >4000 g.

(6) <37 weeks gestation.

(7) 37 or more weeks gestation and <2500 g.

* Percentages are not calculated if <100 records are available for analysis after exclusions.

